

Jiling Cao and Warren B. Moors,\* Department of Mathematics, The University of Auckland, Private Bag 92019, Auckland, New Zealand.  
email: cao@math.auckland.ac.nz and moors@math.auckland.ac.nz

## QUASICONTINUOUS SELECTIONS OF UPPER CONTINUOUS SET-VALUED MAPPINGS

### Abstract

In this paper, we extend a theorem of Matejdes on quasicontinuous selections of upper Baire continuous set-valued mappings from compact (or separable) metric range spaces to regular  $T_1$  range spaces. In addition, we also prove a quasicontinuous selection theorem for a special class of upper semicontinuous set-valued mappings.

### 1 Introduction.

Let  $T : X \rightarrow 2^Y$  be a set-valued mapping with non-empty values. By a *selection*  $f$  of  $T$ , we mean a single-valued mapping  $f : X \rightarrow Y$  such that  $f(x) \in T(x)$  for all  $x \in X$ . A well-known theorem of Michael on selections in [8] claims that any lower semicontinuous set-valued mapping  $T : X \rightarrow 2^Y$  with non-empty closed convex values acting from a paracompact space  $X$  into a Banach space  $Y$  has a continuous selection. However, the conclusion of this theorem fails when lower semicontinuity is replaced by upper semicontinuity. For example, the set-valued mapping  $T : \mathbb{R} \rightarrow 2^{\mathbb{R}}$ , defined by

$$T(x) := \begin{cases} \{1/x\} & \text{if } x \neq 0 \\ \mathbb{R} & \text{if } x = 0 \end{cases}$$

---

Key Words: quasicontinuous, selection, upper Baire continuous  
Mathematical Reviews subject classification: 54C60, 54C65, 26E25  
Received by the editors November 2, 2004

\*The authors are supported by a Marsden fund grant, UOA0422, administrated by the Royal Society of New Zealand

is upper semicontinuous with non-empty closed convex values. Note that this mapping does not even possess a quasicontinuous selection. Recall that a (single-valued) mapping  $f : X \rightarrow Y$  is *quasicontinuous* if for every pair of open sets  $U \subseteq X$  and  $W \subseteq Y$  with  $f(U) \cap W \neq \emptyset$ , there exists a non-empty open set  $V \subseteq U$  such that  $f(V) \subseteq W$ . In a series of papers [4, 5, 6, 7], Matejdes studied the problem of when a set-valued mapping admits a quasicontinuous selection. To achieve his goal, Matejdes introduced the following definition, [4].

**Definition 1.1** ([4]). A set-valued mapping  $T : X \rightarrow 2^Y$  is called *upper Baire continuous* at a point  $x \in X$  if for each pair of open sets  $U$  and  $W$  with  $x \in U$  and  $T(x) \subseteq W$ , there is a subset  $B \subseteq U$  of the second category, having the Baire property, such that  $T(z) \subseteq W$  for all  $z \in B$ .

We shall say that a set-valued mapping  $T : X \rightarrow 2^Y$  is *upper Baire continuous* if it is upper Baire continuous at every point of  $X$ , and a Baire continuous single-valued mapping is just a special case of an upper Baire continuous set-valued mapping. Analogously, one can define lower Baire continuity for a set-valued mapping. However, we shall not do so here, since we are not going to use such a notion in this paper.

The following two facts on (upper) Baire continuity of mappings can be readily proved:

- If  $f : X \rightarrow 2^Y$  is upper Baire continuous with non-empty values, then  $X$  is Baire.
- If a (single-valued) mapping  $f : X \rightarrow Y$  is Baire continuous,  $X$  is Baire and  $Y$  is regular, then  $f$  must be quasicontinuous, [4].

Using the previous two facts Matejdes proved the following theorem.

**Theorem 1.2** ([4]). Let  $X$  be a  $T_1$ -space and  $Y$  be a compact metric space. If  $T : X \rightarrow 2^Y$  is upper Baire continuous with non-empty compact values, then  $T$  admits a quasicontinuous selection.

In [5], it was further shown that the compactness of  $Y$  in the previous theorem can be relaxed to the separability of  $Y$ . The main purpose of this paper is to extend Theorem 1.2 using a different approach. Specifically, in Section 2, we show that the conclusion of Theorem 1.2 still holds when the condition “ $Y$  be a compact (or separable) metric space” is weakened to “ $Y$  be a regular  $T_1$ -space”. The last section is dedicated to the study of quasicontinuous selections of a special class of upper semicontinuous set-valued mappings. Throughout the paper,  $T : X \rightarrow 2^Y$  always denotes a set-valued mapping acting from a topological space  $X$  to a topological space  $Y$  and  $f : X \rightarrow Y$  stands for a

single-valued mapping from  $X$  into  $Y$ . The *graph*  $Gr(T)$  of  $T : X \rightarrow 2^Y$  is defined by

$$Gr(T) := \{(x, y) \in X \times Y : y \in T(x)\}.$$

All of our notation is standard and any undefined concepts may be found in the references.

## 2 An Extension of Theorem 1.2.

Let  $X$  be a topological space. Recall that a set  $A \subseteq X$  is said to be *residual* if  $X \setminus A$  is a set of first category. As usual, the symmetric difference of two sets  $A$  and  $B$  in  $X$  is denoted by  $A\Delta B$ . A set  $A \subseteq X$  is said to have the *Baire property* if  $A\Delta G$  is a set of the first category for some open set  $G \subseteq X$ .

The following characterization for upper Baire continuity of a set-valued mapping is easier to work with than the original definition in Definition 1.1.

**Lemma 2.1.** *A set-valued mapping  $T : X \rightarrow 2^Y$  with non-empty values is upper Baire continuous if, and only if,  $X$  is Baire and for each pair of open subsets  $U$  and  $W$  with  $x \in U$  and  $T(x) \subseteq W$ , there exist a non-empty open set  $V \subseteq U$  and a residual set  $R \subseteq V$  such that  $T(z) \subseteq W$  for all  $z \in R$ .*

PROOF. ( $\Rightarrow$ ). Suppose that  $T : X \rightarrow 2^Y$  is upper Baire continuous. First, by remarks in Section 1,  $X$  must be Baire. Furthermore, by the definition, for each pair of open sets  $U$  and  $W$  with  $x \in U$  and  $T(x) \subseteq W$ , there exists some subset  $B \subseteq U$  of the second category having the Baire property such that  $T(z) \subseteq W$  for all  $z \in B$ . Let  $B = G\Delta C$ , where  $G$  is an open set and  $C$  is a set of the first category. Next, put  $V = G \cap U$  and  $R = G \setminus C$ . Then  $V \subseteq U$  is a non-empty open set and  $R$  is a residual set in  $V$  such that  $T(z) \subseteq W$  for each  $z \in R$ .

( $\Leftarrow$ ). Conversely, suppose that  $X$  is Baire and for each pair of open sets  $U$  and  $W$  with  $x \in U$  and  $T(x) \subseteq W$ , there exists a non-empty open subset  $V \subseteq U$  and a residual subset  $R \subseteq V$  such that  $T(z) \subseteq W$  for all  $z \in R$ . Since  $V$  is of the second category, then  $R$  must be of the second category. In addition,  $R = V\Delta(V \setminus R)$ . Thus,  $R$  has the Baire property as well.  $\square$

Our next theorem extends Theorem 1.2 from a compact (or separable) metric range space to an arbitrary regular  $T_1$  range space.

**Theorem 2.2.** *Let  $X$  be a topological space and  $Y$  be a regular  $T_1$ -space. If  $T : X \rightarrow 2^Y$  is an upper Baire continuous set-valued mapping with non-empty compact values, then  $T$  admits a quasicontinuous selection.*

PROOF. First, by Lemma 2.1,  $X$  must be a Baire space. Let  $\mathcal{M}$  be the family of all upper Baire continuous set-valued mappings from  $X$  to  $Y$  with non-empty compact values such that for every  $H \in \mathcal{M}$ ,  $Gr(H) \subseteq Gr(T)$ . Since  $T \in \mathcal{M}$ ,  $\mathcal{M} \neq \emptyset$ . We define a partial order  $\preceq$  on  $\mathcal{M}$  by writing

$$H_1 \preceq H_2 \text{ if, and only if, } Gr(H_1) \subseteq Gr(H_2).$$

Next, we show that  $\mathcal{M}$  has a minimal element. To this end, let  $\mathcal{M}_0$  be any linearly ordered non-empty subfamily of  $\mathcal{M}$ . Then, define a set-valued mapping  $H_{\mathcal{M}_0} : X \rightarrow 2^Y$  by letting

$$H_{\mathcal{M}_0}(x) := \bigcap \{H(x) : H \in \mathcal{M}_0\}$$

for all  $x \in X$ . Fix an arbitrary point  $x_0 \in X$ . Since  $\{H(x_0) : H \in \mathcal{M}_0\}$  is a linearly ordered family of non-empty compact subsets of  $Y$ ,  $H_{\mathcal{M}_0}(x_0)$  is also a non-empty compact subset of  $Y$ . Now, suppose that  $U \subseteq X$  and  $W \subseteq Y$  are a pair of non-empty open subsets with  $x_0 \in U$  and  $H_{\mathcal{M}_0}(x_0) \subseteq W$ . Then, there must be some element  $H \in \mathcal{M}_0$  such that  $H(x_0) \subseteq W$ . By upper Baire continuity of  $H$  at  $x_0$ , there is a non-empty open set  $V \subseteq U$  and a residual subset  $R \subseteq V$  such that  $H(x) \subseteq W$  for all  $x \in R$ . This implies that  $H_{\mathcal{M}_0}(x) \subseteq W$  for all  $x \in R$ . Thus,  $H_{\mathcal{M}_0} \in \mathcal{M}$ . By Zorn's lemma,  $\mathcal{M}$  has a minimal member, which we will denote by  $\Phi_{\mathcal{M}}$ .

**Claim 1.** *For each pair of open subsets  $U \subseteq X$  and  $W \subseteq Y$  such that  $\Phi_{\mathcal{M}}(U) \cap W \neq \emptyset$ , there exist a non-empty open subset  $V \subseteq U$  and a residual set  $R \subseteq V$  such that  $\Phi_{\mathcal{M}}(x) \subseteq W$  for all  $x \in R$ .*

PROOF. Suppose the contrary. Then, there is a pair of open subsets  $U \subseteq X$  and  $W \subseteq Y$  with  $\Phi_{\mathcal{M}}(U) \cap W \neq \emptyset$  such that for every non-empty open subset  $V \subseteq U$  and every residual subset  $R \subseteq V$  there exists an  $x \in R$  such that  $\Phi_{\mathcal{M}}(x) \not\subseteq W$ . Since  $\Phi_{\mathcal{M}}$  is upper Baire continuous, this implies that  $\Phi_{\mathcal{M}}(x) \not\subseteq W$  for any  $x \in U$ . Next, we define a set-valued mapping  $\Gamma : X \rightarrow 2^Y$  by

$$\Gamma(x) := \begin{cases} \Phi_{\mathcal{M}}(x) \cap (Y \setminus W) & \text{if } x \in U \\ \Phi_{\mathcal{M}}(x) & \text{otherwise.} \end{cases}$$

Then  $\Gamma$  has non-empty compact values. We will show that  $\Gamma$  is upper Baire continuous. Pick any point  $x_0 \in X$ . If  $x_0 \notin U$ , then the result is clear, since  $\Phi_{\mathcal{M}}$  is upper Baire continuous and  $\Gamma \preceq \Phi_{\mathcal{M}}$ . Assume  $x_0 \in U$ . Let  $U'$  and  $W'$  be a pair of open sets with  $x_0 \in U' \subseteq U$  and  $\Gamma(x_0) \subseteq W'$ . Then  $\Phi_{\mathcal{M}}(x_0) \subseteq W \cup W'$ . Thus there exist a non-empty open set  $V' \subseteq U'$  and a residual set  $R' \subseteq V'$  such that  $\Phi_{\mathcal{M}}(x) \subseteq W \cup W'$  for all  $x \in R'$ . Clearly,  $\Gamma(x) \subseteq W'$  for every point  $x \in R'$ . This implies that  $\Gamma$  is upper Baire

continuous at every point of  $U$ . Thus, we have shown that  $\Gamma \in \mathcal{M}$ . But this is impossible since  $\Gamma \preceq \Phi_{\mathcal{M}}$  and  $\Phi \neq \Phi_{\mathcal{M}}$ . Hence we have obtained our desired contradiction.

**Claim 2.**  $\Phi_{\mathcal{M}}$  is single-valued at every point  $x \in X$ .

PROOF. If not, there must exist a point  $x_1 \in X$  such that  $\Phi_{\mathcal{M}}(x_1)$  contains at least two points. Now, pick any point  $y_1 \in \Phi_{\mathcal{M}}(x_1)$ , and then define another set-valued mapping  $\Psi : X \rightarrow 2^Y$  by

$$\Psi(x) := \begin{cases} \{y_1\}, & \text{if } x = x_1, \\ \Phi_{\mathcal{M}}(x), & \text{otherwise.} \end{cases}$$

It is clear that  $\Psi$  has non-empty compact images. Let  $x \in X$  and consider open sets  $U \subseteq X$  and  $W \subseteq Y$  such that  $x \in U$  and  $\Psi(x) \subseteq W$ . By Claim 1, there exist a non-empty open subset  $V \subseteq U$  and a residual subset  $R \subseteq V$  such that  $\Phi_{\mathcal{M}}(x) \subseteq W$  for all  $x \in R$ . It follows that  $\Psi(x) \subseteq W$  for all  $x \in R$ . Thus  $\Psi$  is upper Baire continuous. But,  $\Psi \preceq \Phi_{\mathcal{M}}$  and  $\Psi \neq \Phi_{\mathcal{M}}$ ; which contradicts the minimality of  $\Phi_{\mathcal{M}}$ .

Finally, by Claim 2,  $\Phi_{\mathcal{M}}$  is a Baire continuous selection of  $T$ . Therefore, since  $X$  is Baire and  $Y$  is regular,  $\Phi_{\mathcal{M}}$  is quasicontinuous.  $\square$

### 3 Strongly Injective Set-Valued Mappings

In this section, we shall examine when an upper semicontinuous set-valued mapping acting between topological spaces admits a quasicontinuous selection. Recall that a set-valued mapping  $T : X \rightarrow Y$  from a topological space  $X$  into a topological space  $Y$  is said to be *upper semicontinuous* at a point  $x_0 \in X$  if for every open subset  $V \subseteq Y$  with  $T(x_0) \subseteq V$ , there exists an open subset  $U \subseteq X$  with  $x_0 \in U$  such that  $T(U) \subseteq V$ .

Our considerations are based upon the following notion.

**Definition 3.1.** A set-valued mapping  $T : X \rightarrow 2^Y$  is *strongly injective* if  $T(x_1) \cap T(x_2) = \emptyset$  for any two distinct points  $x_1, x_2 \in X$ .

**Remark 3.2.** If  $f : X \rightarrow Y$  is a surjective mapping, then  $f^{-1} : Y \rightarrow 2^X$  is strongly injective. In particular, the quotient mapping  $q : G \rightarrow G/H$  from a (Hausdorff) group  $G$  onto a coset space  $G/H$  as considered by Michael in [8] is strongly injective. Conversely, for any strongly injective set-valued mapping  $T : Y \rightarrow 2^X$  with non-empty values and  $T(Y) = X$ , it is easy to see that there exists a mapping  $f : X \rightarrow Y$  such that  $T = f^{-1}$ .

Furthermore, we shall also require the definition of property  $(**)$  introduced in [2]. Let  $X$  be a space,  $\mathcal{F}$  a proper filter (or filterbase) in  $X$ . We shall consider the following  $G(\mathcal{F})$ -game played in  $X$  between players  $A$  and  $B$ : Player  $A$  goes first (always!) and chooses a point  $x_1 \in X$ . Player  $B$  responds by choosing a member  $F_1 \in \mathcal{F}$ . Following this, player  $A$  must select another (possibly the same) point  $x_2 \in F_1$  and in turn player  $B$  must again respond to this by choosing a member  $F_2 \in \mathcal{F}$ . Repeating this procedure indefinitely, the players  $A$  and  $B$  produce a sequence  $p := ((x_n, F_n) : n \in \mathbb{N})$  with  $x_{n+1} \in F_n$  for all  $n \in \mathbb{N}$ , called a *play* of the  $G(\mathcal{F})$ -game. We shall say that  $B$  *wins* a play of the  $G(\mathcal{F})$ -game if the sequence  $(x_n : n \in \mathbb{N})$  has a cluster point in  $X$ . Otherwise, the player  $A$  is said to have *won* this play.

We shall call a pair  $(\mathcal{F}, \sigma)$  a  $\sigma$ -filter ( $\sigma$ -filterbase) if  $\mathcal{F}$  is a proper filter (filterbase) in  $X$  and  $\sigma$  is a winning strategy for player  $B$  in the  $G(\mathcal{F})$ -game. Finally, we say that a space  $X$  has *property  $(**)$*  if  $\bigcap \{\overline{F} : F \in \mathcal{F}\} \neq \emptyset$  for each  $\sigma$ -filterbase  $(\mathcal{F}, \sigma)$  in  $X$ . The class of spaces having property  $(**)$  includes all metric spaces [1], all Dieudonné-complete spaces, all function spaces  $C_p(X)$  for compact Hausdorff spaces  $X$ , and all Banach spaces in their weak topologies [2]. Recall that a space  $X$  is a  $q$ -space if for every point  $x \in X$ , there is a sequence  $(U_n : n \in \mathbb{N})$  of neighborhoods of  $x$  such that if  $x_n \in U_n$  for all  $n \in \mathbb{N}$ , the sequence  $(x_n : n \in \mathbb{N})$  has a cluster point in  $X$  (which is not necessarily  $x$  itself). All first countable spaces and all Čech-complete spaces are  $q$ -spaces.

The following theorem may be deduced from [2, Theorem 3.3].

**Theorem 3.3** ([2]). *Let  $T : X \rightarrow 2^Y$  be a strongly injective upper semicontinuous set-valued mapping with non-empty closed values. If  $X$  is a regular  $q$ -space and  $Y$  is a regular space with property  $(**)$ , then for any point  $x_0 \in X$ ,*

$$K := \bigcap_{U \in \mathcal{U}(x_0)} \overline{T(U \setminus \{x_0\})}$$

*is a compact subset of  $T(x_0)$ , where  $\mathcal{U}(x_0)$  is the family of all neighborhoods of  $x_0$  in  $X$  and  $\overline{T(U \setminus \{x_0\})}$  is the closure of  $T(U \setminus \{x_0\})$  in  $Y$ . In addition, the mapping  $T_K : X \rightarrow 2^Y$ , defined by*

$$T_K(x) := \begin{cases} K & \text{if } x = x_0, \\ T(x) & \text{otherwise,} \end{cases}$$

*is upper semicontinuous on  $X$ .*

Note that, in the previous theorem, if  $x_0 \in X$  is not an isolated point, then  $K$  is non-empty.

Our next selection theorem requires the notion of a minimal usco.

**Definition 3.4.** We shall call a set-valued mapping  $\varphi : X \rightarrow 2^Y$  acting between topological spaces  $X$  and  $Y$  an *usco* mapping if for each  $x \in X$ ,  $\varphi(x)$  is a nonempty compact subset of  $Y$  and for each open set  $W$  in  $Y$   $\{x \in X : \varphi(x) \subseteq W\}$  is open in  $X$ . An usco mapping  $\varphi : X \rightarrow 2^Y$  is called a *minimal usco* if its graph does not contain, as a proper subset, the graph of any other usco defined on  $X$ .

**Proposition 3.5** ([3]). *Let  $\varphi : X \rightarrow 2^Y$  be an usco acting between topological spaces  $X$  and  $Y$ . Then  $\varphi$  is a minimal usco if and only if, for each pair of open subsets  $U$  of  $X$  and  $W$  of  $Y$  with  $\varphi(U) \cap W \neq \emptyset$  there exists a non-empty open subset  $V$  of  $U$  such that  $\varphi(V) \subseteq W$ . In particular, every selection of a minimal usco is quasicontinuous.*

**Proposition 3.6** ([3]). *Let  $\varphi : X \rightarrow 2^Y$  be an usco mapping acting from a topological space  $X$  into a Hausdorff topological space  $Y$ . Then there exists a minimal usco  $\psi : X \rightarrow 2^Y$  such that  $\psi(x) \subseteq \varphi(x)$  for all  $x \in X$ .*

**Theorem 3.7.** *Let  $T : X \rightarrow 2^Y$  be a strongly injective upper semicontinuous set-valued mapping with nonempty closed values. If  $X$  is a regular  $q$ -space and  $Y$  is a regular Hausdorff space with property (\*\*), then  $T$  admits a quasicontinuous selection.*

PROOF. For any isolated point  $x \in X$ , pick an arbitrary point  $y_x \in T(x)$ . Next, define the set-valued mapping  $\Phi : X \rightarrow 2^Y$  by,

$$\Phi(x) := \begin{cases} \bigcap_{U \in \mathcal{W}(x)} \overline{T(U \setminus \{x\})} & \text{if } x \text{ is not isolated} \\ \{y_x\} & \text{if } x \text{ is isolated.} \end{cases}$$

By Theorem 3.3 and the subsequent remark,  $\Phi$  has non-empty compact values.

Now, fix an arbitrary point  $x_0 \in X$ . To show that  $\Phi$  is upper semicontinuous at  $x_0$ , we consider two possible cases. If  $x_0$  is an isolated point of  $X$ , then the upper semicontinuity of  $\Phi$  at  $x_0$  is trivial. In the case that  $x_0$  is non-isolated, it follows from the second part of Theorem 3.3. Thus,  $\Phi$  is an usco whose graph is contained in the graph of  $T$ . By Proposition 3.6, there exists a minimal usco  $\psi : X \rightarrow 2^Y$  such that  $\psi(x) \subseteq \Phi(x) \subseteq T(x)$  for all  $x \in X$ . Now, by Proposition 3.5,  $\psi$  has a quasicontinuous selection  $\sigma : X \rightarrow Y$  which in turn is also a selection of  $T$ .  $\square$

**Corollary 3.8.** *Let  $f : X \rightarrow Y$  be a closed mapping from a regular  $T_1$ -space  $X$  with property  $(**)$  onto a regular  $q$ -space  $Y$ . If  $f^{-1}(y)$  is closed for every  $y \in Y$ , then there exists a quasicontinuous mapping  $\varphi : Y \rightarrow X$  such that  $(f \circ \varphi)(y) = y$  for all  $y \in Y$ .*

PROOF. Note that  $f^{-1} : Y \rightarrow 2^X$  is an upper semicontinuous strongly injective set-valued mapping with non-empty closed values. By applying Theorem 3.7,  $f^{-1}$  admits a quasicontinuous selection  $\varphi : Y \rightarrow X$ . Evidently,  $(f \circ \varphi)(y) = y$  for all  $y \in Y$ .  $\square$

**Remark 3.9.** By [2, Theorem 1.2] and an argument similar to that in Theorem 3.7, one can show the following: Let  $T : X \rightarrow 2^Y$  be an upper semicontinuous set-valued mapping from a first countable space  $X$  into a Hausdorff and angelic space  $Y$ . If  $T$  is strongly injective, then it admits a quasicontinuous selection. As a consequence of this result, the condition “ $f^{-1}(y)$  is closed for every  $y \in Y$ ” in Corollary 3.8 can be dropped when  $X$  is Hausdorff and angelic and  $Y$  is first countable; i.e., for any closed mapping  $f : X \rightarrow Y$  from a Hausdorff and angelic space  $X$  onto a first countable space  $Y$ , there exists a quasicontinuous mapping  $\varphi : Y \rightarrow X$  such that  $(f \circ \varphi)(y) = y$  for all  $y \in Y$ .

**Note Added in Proof:** We should observe that the conclusion of Theorem 3.7 remains if we replace the condition “ $T$  is strongly injective” by the weaker hypothesis that “ $T$  is locally strongly injective”; i.e., for each  $x \in X$  there exists a neighborhood  $U$  of  $x$  such that  $T|_U$  is strongly injective on  $U$ .

## References

- [1] J. Cao, *Generalized metric properties and kernels of set-valued maps*, *Topology Appl.*, **146-147** (2005), 603–609.
- [2] J. Cao, W. Moors and I. Reilly, *Topological properties defined by games and their applications*, *Topology Appl.*, **123** (2002), 47–55.
- [3] J. P. R. Christensen, *Theorems of Namioka and R. E. Johnson type for upper semicontinuous and compact valued mappings*, *Proc. Amer. Math. Soc.*, **86** (1982), 649–655.
- [4] M. Matejdes, *Sur les sélecteurs des multifonction*, *Math. Slovaca*, **37** (1987), 111–124.
- [5] M. Matejdes, *On the cliqish, quasicontinuous and measurable selections*, *Math. Bohemica*, **116** (1991), 170–173.



- [6] M. Matejdes, *On the selections of multifunctions*, Math. Bohemica, **118** (1993), 255–260.
- [7] M. Matejdes, *Continuity of multifunctions*, Real Analysis Exchange, **19** (1993/94), 394–413.
- [8] E. Michael, *Continuous selections I*, Ann. Math., **63** (1956), 361–382.

