

LASSO regression used to select important features when predicting wheat yield from various germplasm groups

Muhuddin Rajin Anwar^{a,b,*}, Livinus Emebiri^a, Ryan H. L. Ip^{c,d}, David J. Lockett^b, Yashvir S. Chauhan^e, Ketema T. Zeleke^{b,f}

^aNSW Department of Primary Industries, Wagga Wagga Agricultural Institute, Wagga Wagga, NSW 2650, Australia.

^bGulbali Institute (Agriculture, Water and Environment), Charles Sturt University, Locked Bag 588, Wagga Wagga, NSW 2678, Australia.

^cSchool of Computing and Mathematics, Charles Sturt University, Wagga Wagga, NSW 2650, Australia.

^dDepartment of Mathematical Sciences, Auckland University of Technology, Auckland, New Zealand

^eDepartment of Agriculture and Fisheries (DAF), Kingaroy, Queensland 4610, Australia.

^fSchool of Agricultural, Environmental and Veterinary Sciences, Charles Sturt University, Wagga Wagga, NSW 2650, Australia.

*Corresponding author. *E-mail*: muhuddin.anwar@dpi.nsw.gov.au (M.R. Anwar)

Highlights

- LASSO regression modelling allowed identification of the most influential stress variables affecting the grain yield of bread and Durum wheat grown in field experiments across two sites, multiple years, and two sowing times.
- The influence of crop water supply-demand ratio was an excellent index to summarise water relations and drought stress. Its expression depended upon the growth period examined and the germplasm grouping.
- Evapotranspiration was important only in the vegetative phase (pre-flowering) while drought and heat stress were most important around flowering time.
- Graphs of predicted versus actual yield may help identify wheat genotypes that have a greater-than-expected stress tolerance or avoidance.

Abstract

Bread wheat and Durum wheat genotypes were grown in field experiments at two locations in New South Wales, Australia across several years and using two sowing times ('early' versus 'late'). Genotypes were grouped based on genetic similarity. Grain yield, grain size, soil characteristics, and daily weather data were

36 collected. The Weather data was used to calculate water and heat stress indices for four key growth periods
37 around flowering. LASSO (least absolute shrinkage and selection operator) was used to predict grain yield
38 and to identify the most influential features (a combination of index and growth period). A novel approach
39 involving the crop water supply-demand ratio (SDR) effectively summarized water relations during growth.
40 LASSO predicted grain yield quite well (adjusted R² from 0.57 to 0.98), especially in a set of Durum
41 genotypes. However, the addition of other important variables such as lodging score, disease incidence,
42 weed incidence, and insect damage could improve modelling. Growth period 2 (30 days pre-flowering up
43 to flowering) was the most sensitive for yield loss from heat stress and water stress for most features.
44 Although one group of bread wheat genotypes was more sensitive to water stress (drought) in period 3 (20
45 days pre-flowering to 10 days post-flowering). Evapotranspiration was a significant positive feature but
46 only in the vegetative phase (pre-flowering, period 1). This study confirms the usefulness of statistical
47 modelling as a technique to make predictions that could be used to identify genotypes that are worthy of
48 further investigation by breeders for their stress-tolerance ability.

49
50 **Keywords:** *Triticum aestivum*, LASSO-regression, yield, germplasm, SDR, water-use efficiency, heat
51 stress, water stress

52

53 **1. Introduction**

54 Wheat (*Triticum aestivum* L.) is the world's third most abundant staple cereal food crop, behind maize and
55 rice in terms of production. The OECD/FAO (2023) reports that its annual output for the 2022-2023 period
56 was approximately 800 million tons. In Australia, it is the fifth most exported commodity (OEC, 2023).
57 However, climate change-induced abiotic stresses, such as drought and elevated temperatures, pose
58 significant challenges to wheat production (Collins and Chenu, 2021). These stresses are expanding and
59 intensifying, and impacting global wheat yields (Lobell *et al.*, 2011). Additionally, water scarcity
60 significantly hinders wheat cultivation in Australia, prompting growers, breeders, and agronomists to focus
61 on improving water usage efficiency (Sadras and McDonald, 2012). Durum wheat (*Triticum turgidum* L.
62 subsp. *durum* (Desf.) Husn.) is a secondary wheat crop in Australia – grown for use in pasta production
63 (GRDC GrowNote, 2017). Its annual production in Australia hovers around 0.5 million tonnes – much less
64 than the total for bread wheat of about 36 million tonnes in 2022 (ABS 2024).

65

66 High temperatures during critical crop development stages, such as flowering, can reduce grain yield by
67 directly affecting grain number and grain weight (Trethowan, 2022). This has been shown in several studies,
68 including Stone and Nicolas (1994), Talukder *et al.* (2014), and Wollenweber *et al.* (2003). Even a short
69 period of high temperature during flowering can significantly reduce grain weight and set, especially in
70 sensitive cultivars (Talukder *et al.*, 2013). For example, a field experiment by Nuttall *et al.* (2012) showed

71 that a temperature of 36–38 °C for 6 days after flowering resulted in a 12% reduction in grain number and
72 a 13% loss in grain yield.

73
74 Climate change poses a significant threat to global wheat production. Environmental factors like water
75 shortages and high temperatures significantly impact global wheat production through plant phenotypic and
76 physiological changes (Abhinandan *et al.*, 2018). Studies indicate that for each additional degree of global
77 mean temperature increase, wheat yields could decline by up to 6%, under the assumption of no CO₂
78 fertilization, continued effective management practices, and no changes in crop genetics (Asseng *et al.*,
79 2015; Zhao *et al.*, 2017). This impact is already evident in Australia, where simulations suggest a huge and
80 concerning 27% decline in water-limited potential wheat yield from 1990 to 2015 (Hochman *et al.*, 2017).
81 This decrease is likely attributable to a combination of stressors such as seasonal rainfall and increased
82 temperatures, coupled with the limited ability of increased atmospheric concentration (CO₂) to fully
83 compensate for these negative factors (Wang *et al.*, 2017; Li *et al.*, 2022). While climate change and climate
84 variability present major hurdles, analysing the connections between climate, soil, and wheat yield
85 empowers us to design actionable strategies to mitigate yield losses. By unravelling the relative importance
86 of various variables, we can guide future research towards developing climate-resilient wheat cultivars and
87 innovative management practices, ultimately transforming vulnerability into opportunity.

88
89 In any individual wheat crop, in addition to gross climatic and edaphic influences, many other biotic and
90 abiotic stresses will affect crop growth and yield. In this work, we were particularly interested in abiotic soil
91 water deficit and heat stresses, especially during the reproductive period of the crop's growth. We wanted
92 to explore how we can use soil characteristics and weather variables to improve the prediction of phenology
93 and yield via modelling. In addition, we wanted to identify the most influential variables and the most
94 sensitive crop growth period upon which these variables act. The least absolute shrinkage and selection
95 method (LASSO) has shown good utility for identifying the most influential variable in a multiple-
96 regression situation of this type (Tibshirani, 1996; Didari *et al.*, 2023). This methodology may assist wheat
97 breeders by identifying different influential variables depending on the type of wheat germplasm being
98 examined. In addition, if other traits are of interest, the same approach can be used to dissect the influential
99 variables (Shafiee *et al.* 2021).

100
101 The specific objectives of this study were as follows.
102 Firstly, to utilise an existing wheat dataset comprising six sets of bread wheat and durum wheat germplasm
103 (each consisting of a variable numbers of genotypes), grown in experiments at two sites and up to two
104 sowing times over several years, for research into the possibility of predicting grain yield from a suite of

105 weather- and soil-based climatic variables: particularly, crop water-use, crop water stress, and crop heat
106 stress.

107
108 Secondly, to determine which of those variables were most influential and effective for predicting yield
109 using the relatively new regression technique of LASSO, and to see whether the germplasm groups showed
110 different responses.

111
112 Thirdly, to calculate the weather and soil-based variables across four different crop growth periods
113 (overlapping development stages), and to see which weather and soil-based variables were most influential
114 in predicting yield.

115
116 Fourthly, using the combined findings, to identify the most critical growth period for crop damage (yield
117 loss) from water stress and/or heat stress in the different germplasm groups, with the aim of informing wheat
118 breeders which crop traits merit attention for reducing losses of potential yield from water- and heat-stress.

119
120

121 **2. Materials and methods**

122
123 **2.1 Study area and soil data**

124 Two typical rainfed crop-livestock growing locations (Leeton and Wagga Wagga) in southeast Australia,
125 encompassing different climatic conditions, were selected for analysis (Table 1) utilising an existing dataset
126 previously published for other purposes (Sissons *et al.*, 2018; Zeleke *et al.*, 2023). The soils across the sites
127 are predominantly Wunnamurra Clay (Leeton) and kandosol (Wagga Wagga) according to the Australian
128 Soil Classification (Isbell and National Committee on Soil and Terrain, 2021). The soil properties at these
129 sites have been summarized by others (Xing *et al.*, 2017; Wang *et al.*, 2017). Briefly, at Leeton, the plant
130 available water holding capacity (PAWC) was 293mm, to a total soil depth of 1.8m, pH (1:5 water) ranged
131 from 7.2 – 8.9, bulk density (g/cc) of 1.20 to 1.40 and initial nitrate (NO₃) was 81 kg/ha. At Wagga Wagga,
132 the PAWC was 128mm, to a total soil depth of 1.25m, pH (1:5 water) ranged from 6.2 – 6.9, bulk density
133 was 1.37 to 1.56 g/cc, and initial NO₃ was 69 kg/ha. These sites have been the subject of parameter
134 verification in wheat cropping systems (Anwar *et al.*, 2015; Anwar *et al.*, 2022) especially for the numerous
135 initial values and parameters required for running the APSIM crop growth model (<https://www.apsim.info/>).

136

137 **2.2 Field experiments and agronomy**

138 The layout of the field experiments, the details of the genotypes used, and the agronomy used during each
139 year are detailed in a previous publication (Sissons *et al.*, 2018; Zeleke *et al.*, 2023). Briefly, the experiments
140 were conducted at Leeton in 2011 and 2015 and at Wagga Wagga in 2012, 2018 and 2019. Two sowing
141 times were used at each site/year: ‘early’ and ‘late’, in order to maximise the differences in the weather
142 experienced by the crops. Not all germplasm sets were grown in every site/year. Here we note, in addition,
143 that some lodging occurred in the field along with some fungal disease; the genotypes were variously
144 affected but the scoring used was unfortunately inconsistent. Consequently, in this analysis these factors
145 were not included in the LASSO modelling (see below) and may have contributed to some imprecision in
146 the predicted values.

147

148 **2.3. Wheat Germplasm groups**

149 **ABD lines:** These are advanced breeding lines of wheat (*Triticum aestivum*) produced at the International
150 Maize and Wheat Improvement Centre (CIMMYT), Mexico. They are comprised of the line selections from
151 the high-temperature wheat yield trials (HTWYT), and selections made for their large grain size (Sissons *et*
152 *al.* submitted). Hereafter referred to as “BreadWheat_ABDLines”.

153 **Elite wheat:** These are bread wheat varieties of historical significance, recently released cultivars, and
154 parents used in breeding programs by the major private breeding companies (InterGrain, LongReach, and
155 Australian Grain Technologies) in Australia. These varieties have been bred to meet the specific needs of
156 Australian growers, such as resistance to diseases and pests, tolerance to heat and drought, good grain quality,
157 and high yield, and were chosen based on being potentially heat tolerant (or in some cases intolerant)
158 according to Australian breeder recommendations and the literature (Sissons *et al.*, submitted). Hereafter
159 referred to as “BreadWheat_Elite”.

160 **Landrace wheat:** The bread wheat landraces were sourced from heat-prone areas in Afghanistan, Iran, Iraq,
161 and India. They were identified using FIGS (focused identification of germplasm strategy), an approach that
162 uses environmental parameters described in plant germplasm collection sites as selection criteria to identify
163 materials that most likely have undergone selection pressures for the target parameters (Sissons *et al.*, 2018).
164 Hereafter referred to as “BreadWheat_Landraces”.

165 **Tamaroi x Saintly durum bi-parent population:** These are durum wheat doubled-haploids, which were
166 produced from F₁ plants of a cross between the SA-bred variety, Saintly and the NSW-bred variety, Tamaroi.
167 Saintly has a reputation for performing well in seasons with terminal drought stress, while the variety

168 Tamaroi has a very high inherent 1000-kernel weight but is susceptible to heat stress. Hereafter referred to
169 as “Durum_Biparent”.

170 **Durum elite:** The durum wheat germplasm comprised a worldwide collection trialled for heat tolerance in
171 southern Australia (Collins *et al.*, 2017). They included commercial durum varieties and breeding lines,
172 along with tetraploid wheat landraces sourced from heat-prone regions by using the FIG strategy (Street *et*
173 *al.*, 2016). They have been shown to exhibit significant variability for tolerance/intolerance to late-sown
174 heat stress (Sissons *et al.*, 2018) and natural heat waves (Emebiri *et al.*, 2024). Hereafter referred to as
175 “Durum_Elite”.

176 **Bread wheat NI lines:** The near-isogenic (NI) lines were created from a cross of wheat varieties Drysdale
177 and Waagan. Both parents are semi-dwarf varieties and carry genetic loci for intolerance and tolerance,
178 respectively, to both booting and grain filling stage heat stress (Shirdelmoghanloo *et al.*, 2016; Erena *et al.*,
179 2021). The NI lines were created by using molecular markers to identify single Drysdale × Waagan F_{2:8}
180 plants that were heterozygous for genetic loci located on wheat chromosomes 2B, 3B and 6B; then the
181 progeny of these plants was screened to identify plants homozygous for each allele at the respective loci
182 (Erena *et al.*, 2021). Hereafter referred to as “BreadWheat_NILines”.

183

184 **2.4. Soil water balance**

185 Rainfall, evapotranspiration (ET), runoff, and drainage are key factors that affect how much water is
186 available to crops (Unkovich *et al.*, 2018; Unkovich *et al.*, 2023). In this study, temperature, rainfall,
187 simulated initial soil water content, and simulated soil water content at harvest, were used to calculate the
188 soil water balance following the procedure of He and Wang (2019). We used the pre-validated Agricultural
189 Production Systems sIMulator (APSIM) that simulates the key biophysical processes related to crop growth
190 and production, water, carbon, and N cycling in the soil-plant system (Holzworth *et al.*, 2014). Published
191 studies have also used the APSIM model to calculate hydraulic parameters for wheat cropping systems (such
192 as, soil water content at sowing and harvest, water use, runoff, drainage, and soil evaporation). The
193 parameters in the Soil Water module of APSIM were the same for our sites as those used in other published
194 work (Liu *et al.*, 2014; Zeleke and Nendel, 2019; Wang *et al.*, 2017; Xing *et al.*, 2017). To estimate the
195 initial soil water content at the start of the experimental period (2011), we assumed that the starting soil
196 water on 1 January 2002 was equal to LL15 (water content at 15 bar suction). Then by running APSIM for
197 the 2002-2011 period using actual weather data we simulated the initial soil water in 2011 (He and Wang,

2019). The LL15 value was determined in the Wagga Wagga Agricultural Institute soil moisture analysis laboratory (Anwar *et al.*, 2022). The APSIM model was then run continuously until the end of the experimental period (31 December 2019), without resetting soil water conditions, to obtain the “initial soil water at sowing” and the “soil water at harvest” for each of the wheat experiments (Sissons *et al.*, 2018). The APSIM crop sequence used in the 10-year run-up period before the wheat experiments commenced was a typical one used in the wheat growing regions in Australia: wheat(W)-canola(C)-chickpea (CP)-W-C-CP-W-C-CP-W.

Total crop water use (WU) expressed as evapotranspiration (ET) was calculated by subtracting the final soil water content at harvest from the initial soil water content at sowing and adding the amount of irrigation and rainfall received during the growing season (1).

209

$$210 \quad ET = P + SW_s - SW_h - R - D \quad (1)$$

211

212 where P, R and D are cumulative rainfall, runoff, and deep drainage from the day of sowing to harvest, and
213 SW_s and SW_h are soil water at the sowing and harvest dates, respectively (Yang *et al.*, 2016).

214

215 In contrast, Transpiration (T), which does not include soil evaporation (E) (eqn 1), was calculated using the
216 following soil water balance equation (2) (Yang *et al.*, 2016).

217

$$218 \quad T = P + SW_s - SW_h - R - D - E \quad (2)$$

219

220 The APSIM soil water module also calculates daily potential evapotranspiration using the Priestley-Taylor
221 method (APSIM, 2023), which is based on the physiological relationship between crop yield and
222 evapotranspiration (Paredes *et al.*, 2014; Trout and DeJonge, 2017; Akumaga and Alderman, 2019).

223

224 **2.5. Water supply-demand ratio (SDR)**

225 The APSIM model calculates a water-deficit index (Chapman *et al.*, 1993; Chenu *et al.*, 2011), also known
 226 as the "water supply" and "water demand" ratio, which indicates how well the water extractable by the
 227 crop's roots (water supply) meets the crop's potential transpiration (water demand). The crop water supply
 228 is calculated for each layer of the soil where roots are present and depends on the root growth and soil
 229 property of each layer. The water demand is the amount of water the crop would have transpired in the
 230 absence of soil water constraint. It is estimated daily based on the amount of crop growth on that day and
 231 the atmospheric saturation vapor pressure deficit.

232
 233 Water supply-demand ratio (SDR) is the ratio between water supply and water demand, bounded between
 234 0 and 1, which indicates if the plant is water-stressed.

$$235 \quad SDR = \begin{cases} \min\left(\frac{Supply}{Demand}, 1\right), & Demand > 0 \\ 1, & Demand = 0 \end{cases} \quad (3)$$

237 When $SDR = 1$, there is no water stress. Otherwise, the plant is stressed. Based on SDR, we define water
 238 deficiency (D) such that:

$$239 \quad D = 1 - SDR \quad (4)$$

240 The interpretation is the opposite of SDR. When $D = 0$, there is no water stress. Positive D indicates stress.
 241 Daily deficiency values were calculated and were accumulated within the following four crop development
 242 periods, each spanning approximately 30 days (see below).

243
 244 **2.6. Wheat developmental period**

245 Abiotic stress during the reproductive stage of plants (anthesis and grain filling) has a significant effect on
 246 grain yield and quality. The critical period for abiotic stress is the time when plants are most sensitive to
 247 these stresses. Some previous studies have defined the critical period as 30 or 45 days before to 0 days after
 248 50% anthesis (Fischer, 1985). Other studies have found that the critical period is narrower, spanning only
 249 about 20 days before to 10 days after anthesis (Ortiz-Monasterio *et al.*, 1994; Abbate *et al.*, 1995). More
 250 recently, Slafer *et al.* (2023) found that the critical period for wheat is from 30 days before to 10 days after
 251 anthesis.

252 In this study, we defined and examined four contrasting crop growth periods based on previously published
253 studies:

254

255 **Period 1:** From sowing to the day of flowering (varying lengths)

256 **Period 2:** From 30 days before flowering to the day of flowering (30 days total)

257 **Period 3:** From 20 days before flowering to 10 days after flowering (30 days total)

258 **Period 4:** From 15 days before flowering to 15 days after flowering (30 days total)

259

260 We chose these 30-day intervals based on the findings of previous studies (Fischer, 1985; Slafer *et al.*, 2023).
261 There is considerable chronological overlap between these periods, but we wanted to test which of these
262 periods might be most sensitive to stress effects on grain yield. By definition, Period 2 overlaps with Period
263 3 by 67%; Period 2 overlaps with Period 4 by 50%; and Period 4 overlaps with Period 3 by 83%. The degree
264 to which Period 1 overlaps with the others depends on the interval from sowing to flowering (in days). The
265 means and ranges for the sowing-to-flowering interval for each germplasm group across each
266 Site/Year/Sowing-time combination are given in Table S2 in the Supplementary Material. The overall mean
267 of this duration was 106.0 days. The mean overlap (and the ranges) between Period 1 and Period 2 was 28.8%
268 (21.5% - 38.8%). For periods 1 and 3 the corresponding data were 19.2% (14.3% – 25.9%). For periods 1
269 and 4 they were 14.4% (10.8% – 19.4%).

270

271 **2.7. Statistical techniques and least absolute shrinkage and selection operator (LASSO)**

272 Firstly, we examined some basic summary statistics for each germplasm set and each sowing time (across
273 both sites and all years). Since the “early” (coded as “1”) and “late” (coded as “2”) sowing times were
274 designed to present the crops with contrasting stress environments, we were expecting to see quite large
275 differences in means and ranges for the traits of interest.

276

277 Secondly, to investigate the impact of daily abiotic stress indices (heat stress, water deficit and
278 evapotranspiration) on wheat yield, accumulated over four key growth stages, we used the following
279 approach.

280
281 Pearson correlation coefficients were calculated to assess the relationship between yield, 1000 grain weight
282 (TGW), evapotranspiration (ET), transpiration (Tran), accumulated water deficit (water supply-demand
283 ratio), and heat stress (number of days with temperatures $>30^{\circ}\text{C}$). Correlation analysis was restricted to
284 Period 3 only (20 days before flowering to 10 days after flowering, see above) because this flowering period
285 has proven to be the most important with respect to yield in other published papers (see above). All data
286 were normalized to zero mean and unit variance prior to analysis.

287 Thirdly, to study the relationship between wheat yield (the target trait) and daily stress indices (the
288 explanatory variables) accumulated over critical periods of growth (the four “periods”) and across different
289 sets of genotypes (the germplasm groups) we undertook LASSO regression analysis. In linear models such
290 as multiple linear regression models, it is often assumed that the explanatory variables are independent
291 (Monahan, 2011). When explanatory variables are correlated, multicollinearity is said to exist (Kutner *et al.*
292 2005). As a result of multicollinearity, the estimation of coefficients can become unstable, leading to
293 unreliable estimates. In some extreme cases, the regression coefficients do not reflect the inherent
294 relationship between the explanatory variable and the response variable. For example, a negative coefficient
295 may be obtained although the relationship should be positive.

296 For better interpretability, many statistical methods have been proposed to deal with multicollinearity, many
297 of which are aimed at minimising the prediction error while forcing (i.e., “shrinking”) some of the regression
298 coefficients to zero, hence effectively removing some of the explanatory variables and highlighting the most
299 influential ones (Dormann *et al.*, 2013). Among these methods, LASSO (least absolute shrinkage and
300 selection operator; Tibshirani, 1996) is a popular choice. In this work, we adopted LASSO to find the best
301 subset of explanatory variables from the large initial number. To obtain scientifically sensible regression
302 coefficients, constraints were imposed on them in the estimation procedure. Specifically, the coefficients of
303 variables related to heat and water stresses were set to be non-positive. The computations were performed
304 using the “glmnet” package in R (Friedman *et al.*, 2010).

305 For data preparation, summarisation, and graphics extensive use was made of the “tidyverse” R packages
306 (Wickham *et al.*, 2019), the RStudio GUI (RStudio Team 2023), and the R software suite (R Core Team,
307 2023).

308

309 **3. Results**

310 **3.1. Data summaries across sites, genotype groups and sowing time**

311 The Wagga Wagga soil, compared to Leeton, is a shallower and more dense soil, with lower pH, which
312 holds much less water than Leeton (Table 1). Both sites face sizeable year-to-year variations in the climate
313 variables (rain, solar, and temperatures). Wagga Wagga gets more rain, both overall and during the growing
314 season, with Wagga Wagga's temperatures being slightly cooler than Leeton.

315 We note that the number of genotypes is not the same between the two sowing times within a germplasm
316 group (category), although there was considerable overlap. The frequencies of concurrence of genotypes
317 across “site_year_sowing-time” are given in Table S1. This was due to practical issues, such as the lack of
318 seed supply. The “BreadWheat_NILines” group was only sown once, while the “Durum_Elite” category
319 had only a small number of genotypes. Both of these groups were excluded from the LASSO analysis.

320 While there was a large variation within each germplasm category, the grain yield (Table 2) was always
321 substantially lower in sowing-time_2 due to higher stress levels with an overall range of nearly 9 tonnes per
322 hectare to less than 0.2 tonnes per hectare. Grain size was similarly reduced in sowing-time_2 except for
323 the “Durum_Biparent” category (Table 2).

324 The mean total transpiration (Tran) and mean total evapotranspiration (ET) were always higher in sowing-
325 time_2 (Table 3) due to the crops growing in a hotter and drier period of the year, with ET always greater
326 than Tran (as expected). Water use efficiency (both WUE_TRAN and WUE_ET) were much reduced in
327 sowing-time_2 compared to sowing-time_1, often by more than 50%. The WUE_Tran ranged overall for
328 individual genotypes from 57.5 to 0.92 kilograms of grain per hectare per millimetre, whereas WUE_ETA
329 ranged from 33.5 to 0.6 kilograms of grain per hectare per millimetre (Table 3).

330

331 **3.2. Correlation analysis**

332 Tables 4 to 9 shows the correlation coefficients between yield, 1000 grain weight (TGW),
333 evapotranspiration (ET), transpiration (Tran), accumulated water deficit (SDR = supply-demand ratio), and
334 heat stress ($H>30$ = number of days with temperatures >30 °C). Generally, the correlations between traits
335 were highly significant (either positively or negatively) within each germplasm category but significance
336 levels were much lower (or non-existent) in the “BreadWheat_Landraces” and the two Durum categories.
337 The Tran and ET variables were always highly positively correlated (as expected). The $H>30$ (index of heat
338 stress) was usually highly positively correlated with both Tran and ET but was not significant in the
339 “BreadWheat_Landraces” category (Table 6) nor for ET in the “Durum_Elite” category (although the
340 number of values was small, $n = 10$, Table 9).

341 The TGW and grain yield were generally positively correlated but again not within the
342 “BreadWheat_Landraces” group (Table 6), and were strongly negative in the “Durum_Biparent” material
343 (Table 8). The SDR (index of water stress) was usually negatively correlated (when significant) with the
344 other traits but a contrasting positive correlation was seen in “BreadWheat_ABDLines” material with TGW
345 (Table 4), and with Tran and ET in “Durum_Biparent” material (Table 8).

346 Tran and ET were generally significantly negatively correlated with both yield and TGW, except in the
347 “BreadWheat_Landraces” group (Table 6) and in “Durum_Biparent” (Table 8). As expected, the numerous
348 genotypes in the “BreadWheat_Landraces” category provided the greatest range in performance (yield and
349 TGW), water use (Tran and ET), and water use efficiency (WUE_Tran and WUE_ET), plus less rigid inter-
350 trait correlations.

351

352 **3.3. Distributional characteristics of water and heat stress and evapotranspiration**

353 Environmental factors like water shortages and high temperatures significantly impact global wheat
354 production through plant phenotypic and physiological changes (Abhinandan *et al.*, 2018). For each of the
355 four growing periods in this study, the intensity and frequency of water stress (calculated via supply-demand
356 ratio; SDR) and heat stress including evapotranspiration are summarised below for individual genotypes
357 within six germplasm groups.

358

359 In Figure 1, each boxplot shows the distribution of accumulated values of SDR for a single germplasm
360 group and growing period combination. There is a considerable amount of variability in SDR within each
361 germplasm group and growing period. The boxes, which represent the interquartile range (IQR), span a

362 wide range of values in most cases. The whiskers, which extend to the most extreme data points not
363 considered outliers, also show a wide range of values for many of the groups and periods. The medians
364 (represented by the horizontal lines within the boxes) are generally lower for groups and periods with higher
365 SDR. For example, in the first growing period, the median SDR for the BreadWheat_Elite group is around
366 10, while the median SDR for the same group at third growing period decreased to about 4.8. The dispersion,
367 or spread, of the data is also influenced by SDR. The boxes tend to be wider for groups and periods with
368 higher SDR, indicating that there is a greater range of SDR values within those groups. For example, in the
369 second growing period, the box for the BreadWheat_Elite group is wider than the third growing period. In
370 contrast, BreadWheat_NILines group didn't show dispersion but higher SDR values in the first and fourth
371 growing periods compared to the second and third growing periods. From Figure 1, it can be seen that the
372 SDR decreases from growing period 1 to growing period 4 for all the genotypes. For three of the genotypes
373 (BreadWheat_ABDLines, BreadWheatNILines, Durum_Wheat), the lowest SDR is in growing period 2.

374

375 The potential evapotranspiration (Fig. 2) exhibits considerable variability within each germplasm group and
376 growing period, as evidenced by the interquartile ranges and whiskers of the boxes. The degree of dispersion
377 in evapotranspiration values differs across germplasm groups and growing periods. For instance,
378 Durum_Elite lines generally demonstrate a more compact distribution of evapotranspiration values
379 compared to BreadWheat_ABDLines, suggesting greater consistency in water use within the Durum_Elite
380 group. The median evapotranspiration values vary across germplasm groups and growing periods. Notable
381 trends include BreadWheat_ABDLines tend to have higher median evapotranspiration values than other
382 groups across most growing periods. Durum Elite lines generally exhibit lower median evapotranspiration
383 values, particularly in growing periods 2 and 3. BreadWheat_Landraces display a wider range of median
384 evapotranspiration values across growing periods. Th Potential evapotranspiration of growing period 1 > 4 >
385 3 > 2. Growing period 1 has the highest potential evapotranspiration and growing period 2 has the lowest
386 potential evapotranspiration. Compared to the other genotypes, BreadWheat_Landraces has the highest
387 potential evapotranspiration for each of the respective growing periods and high variability in
388 evapotranspiration across growing periods, suggesting that water use strategies of genotypes may vary
389 depending on environmental conditions and crop developmental stages.

390

391 Figure 3 shows a wide range of variability in heat stress within each germplasm group and growing period.
392 The boxes show the middle 50% of the data, with the whiskers extending to the 10th and 90th percentiles.

393 For example, in the BreadWheat_ABDLines group, the heat stress ranges from 0 to 15 days across the
394 growing period. The median heat stress is also different for each germplasm group and growing period. For
395 example, the median heat stress for the BreadWheat_ABDLines group is about 7 days in the third growing
396 period, while the median heat stress for the Durum_Biparent group is higher (about 15 days) in the same
397 growing period. Figure 3 show that heat stress growing period $1 < 2 < 3 < 4$. Growing period 1 has the
398 lowest heat stress and growing period 4 has the lowest potential evapotranspiration. Durum_Biparent had
399 the highest stress for a given growing period compared to the other genotypes.

400

401 3.4. LASSO feature selection

402 Our wheat data consisted of six germplasm categories (Table S2); however, when fitting a LASSO model,
403 specific criteria must be met. In our case, the “BreadWheat_NILines” category only had one sowing time
404 at one location; hence there was no variation in the explanatory variables, and this category was excluded
405 from the final modelling. Similarly, the “Durum_Elite” category had too few observations (2 genotypes
406 only) to allow the fitting of the explanatory variables. The interpretation of coefficients from LASSO is
407 almost the same as in multiple regression models. The only difference is that LASSO ‘forces’ some of the
408 coefficients to zero. Table 10 shows the estimated coefficients from LASSO and overall model performance.

409

410 In all four major germplasm categories, the effect of ET on yield was effectively zero except in Period 1
411 (where it presumably influenced vegetative biomass, which led to more yield), and in Period 4 for
412 “Durum_Biparent” genotypes. Notably, in Period 1, the effect of ET on yield was higher for
413 “BreadWheat_ABDLines” and “BreadWheat_Elite” compared to the other two germplasm groups. Heat
414 stress (H) was damaging in all periods for the first two bread wheat categories and the “Durum_Biparent”
415 set, but less so for the “BreadWheat_Landraces” set in Period 3. Yet, heat stress in Period 2 was found to
416 be highly damaging for the “BreadWheat_Landraces” group. The results were more mixed for the water
417 stress index (D), particularly detrimental in “BreadWheat_ABDLines” and in Period 3.

418

419 For BreadWheat_ABDLines (Table 10), wheat grain yield was found to be most severely affected by water
420 stress in period 3, followed by heat stress in period 1. For each unit increase in water stress in period 3, yield
421 is expected to decrease by 0.789 t/ha, assuming all other factors remain unchanged. Water stress in period
422 4, and ET in periods 2 to 4, were found to be relatively less influential to the grain yield. In

423 "BreadWheat_Landraces" germplasm category, heat stress during period 2 had the strongest negative impact
424 on wheat grain yield, reducing it by 0.725 t/ha. In contrast, water stress and evapotranspiration in all periods
425 had minimal to no effect on grain yield. Among the germplasm categories in period 2, BreadWheat_Elite
426 experienced the greatest yield reduction due to water stress, with an expected decrease of 0.501 t/ha and
427 heat stress followed closely (yield decline of about 0.449 t/ha). Notably, evapotranspiration had no impact
428 on grain yield for BreadWheat_Elite in periods 2, 3, and 4. Durum_Biparent appears to be less sensitive to
429 water stress and heat stress than BreadWheat_ABDLine. In Durum_Biparent, heat stress had the greatest
430 impact in period 2, with an expected yield decrease of 0.361 t/ha. This was followed by water stress with a
431 decrease of 0.241 t/ha. Evapotranspiration had a positive effect on grain yield in periods 1 and 4, with
432 increases ranging from 0.224 to 0.278 t/ha. However, it had no impact on yield in periods 2 and 3.

433

434 **3.5. Yield prediction**

435 LASSAO modelling predicts yield reasonably well (Figure 4) with highly significant positive regression
436 between observed and predicted values: the "Durum_Biparent" relationship being particularly strong. There
437 are some outlying groups of genotypes, for example in the "BreadWheat_Elite" category but these were
438 very low yielding genotypes. As shown in Table 4, the root mean squared errors ranged from 0.119 to 0.976
439 t/ha across the four genotypes and the adjusted R^2 ranged from 0.57 to 0.98. So, overall, the LASSO
440 approach is working well at predicting crop outcomes, especially for "Durum_Biparent", from weather-
441 based and soil-based indices. Some other explanatory variables (not considered in this study) are required
442 to improve further the goodness of fit, such as disease scores, lodging scores, weed measurements, and crop
443 plant density.

444

445 **4. Discussion**

446 Climate change throws a complex web of challenges at crop production, weaving together water deficits,
447 scorching heat, and fluctuating evaporative demands (Anwar *et al.*, 2015; Kerr *et al.*, 2022). These
448 interwoven environmental stresses act like a multi-pronged attack, inflicting far more damage on plant
449 growth and yield than individual stressors do in isolation (Pandey *et al.*, 2017). This "synergistic effect" can
450 significantly cripple crop production, exceeding initial projections, as evidenced by numerous studies
451 (Mittler, 2006; Prasad *et al.*, 2011).

452

453 This study investigated the combined effects of abiotic stresses: heat, water deficit (SDR), and
454 evapotranspiration, on various wheat germplasm categories in Australia. The findings highlight the intricate
455 and multifaceted nature of understanding how multiple stressors impact crop performance.

456

457 Delayed sowing results in longer crop emergence time, slower growth, less ground cover, lower biomass,
458 higher non-productive (evaporation) component of water balance. Late sown crop is exposed to higher
459 temperature and evapotranspiration (ET) during critical crop development stage. Early sown crop has a
460 deeper rooting system to access subsoil water during the reproductive growth stage (Zelege and Nendel,
461 2019). For all the growth periods considered in this study (periods 1 to 4), the correlation between
462 explanatory variables (TGW [1000 grain weight], ET, Tran [transpiration], SDR, H>30 [number of days
463 with temperatures >30 °C]) and dependent variable (grain yield) is different for different germplasm groups
464 (results only shown for Period 3). This can be due to the inherent difference of the genotypes or due to
465 pooled data from two sites and two sowing times. Heat, evapotranspiration, and transpiration are negatively
466 correlated with yield. One would expect that the more a crop transpires, the higher will be the yield. However,
467 in our data higher rainfall (or higher ET or Tran) years were affected by lodging, resulting in lower yield.

468

469 Our research confirms that climate change presents significant challenges for wheat production. The
470 different growing periods exhibited variations in water stress, evapotranspiration, and heat stress (H>30
471 days), demonstrating the potential for diverse climatic pressures throughout the growing season (Nuttall *et al.*, 2018). These stresses were found to significantly impact grain yield and plant characteristics like
472 thousand-grain weight.
473

474

475 Interestingly, this study emphasizes that the combined effect of these stressors isn't simply additive.
476 Interactions between factors like heat and water deficit can be complex and vary depending on the specific
477 germplasm category and growing period. For example, while heat stress generally reduced yield in most
478 wheat germplasm categories tested here, its impact was less pronounced in the "BreadWheat_Landraces"
479 group in period 3, while water stress in period 2 had the largest detrimental effect for this group. Conversely,
480 the "Durum_Biparent" group seemed less sensitive to stress overall, even showing a positive response to
481 increased evapotranspiration in some periods (Sinha *et al.*, 2021; Ru *et al.*, 2023).

482 These findings underline the need for nuanced approaches to managing wheat crops under increasing
483 climate variability (FAO, 2016). Selecting stress-tolerant varieties and implementing targeted strategies
484 based on specific environmental conditions and germplasm characteristics will be crucial for ensuring food
485 security in a changing climate. Further research exploring additional stress factors and their interactions will
486 also be vital for optimizing wheat production and resilience.

487
488 While the LASSO model effectively captured the main stress effects (Shafiee et al. 2021), it's important to
489 acknowledge the limitations. The observed stress-yield relationships likely involve intricate interactions that
490 the model might not fully capture. For instance, the contrasting response of "BreadWheat_Landraces" to
491 heat stress across different periods suggests potential moderating factors or complex physiological
492 mechanisms at play. Further research delving deeper into these interactions and incorporating additional
493 stress factors like salinity or nutrient deficiency could provide a more comprehensive understanding of how
494 multiple stresses collectively impact wheat performance (Teixeira *et al.*, 2013; Ru *et al.*, 2023).

495
496 Despite these known limitations, the LASSO model demonstrated promising results in predicting yield
497 based on weather and soil-based indices, particularly for the "Durum_Biparent" group. This highlights it's
498 potential as a tool for:

499 1) Identifying stress-tolerant genotypes: by analysing the LASSO coefficients and stress responses
500 across diverse germplasm, researchers can prioritize genotypes with inherent resistance or resilience to
501 specific stress combinations. This can be achieved by identifying germplasm categories that exhibit
502 consistently lower yield reductions under various stress combinations.

503 2) Targeted stress mitigation strategies: understanding which stress factors are most critical for specific
504 genotypes and growth periods allows for tailored interventions. For example, if water stress is the primary
505 limiting factor for a particular germplasm category during a specific growth period, implementing irrigation
506 scheduling strategies can be crucial. Conversely, for germplasm categories sensitive to heat stress, exploring
507 heat stress management techniques such as by earlier sowing or by growing earlier maturing varieties or
508 breeding for heat tolerance can be prioritized.

509

510 To summarise, the LASSO analysis provided valuable insights into the diverse and complex ways that
511 abiotic stresses impact wheat yield across different germplasm categories. While further research is needed
512 to fully understand the intricate interactions between stresses, this study demonstrates the potential of
513 LASSO as a tool for predicting and managing stress impacts, ultimately contributing to improved wheat
514 production and food security in a changing climate.

515

516 **5. Conclusion**

517 In this study we demonstrated how LASSO can be used to identify bread wheat and Durum wheat genotypes
518 with stress-tolerance ability within germplasm groupings using data from multi-site and multi-year field
519 experiments grown in NSW Australia. Grain yield, soil characteristics, and daily weather data were recorded
520 to predict grain yield using stress indices. LASSO predicted grain yield well but adding other variables like
521 lodging score, disease incidence, weed incidence, and insect damage could improve this technique. Not all
522 growing periods were predicted well. We found that the growth period 30 days pre-flowering up to flowering
523 was sensitive for yield loss from heat and water stress as compared to other three periods of similar duration.
524 The study confirms the usefulness of statistical modelling in identifying genotypes worthy of investigation
525 by breeders.

526

527 **Acknowledgment.** We acknowledge the contributions of Dr. Nicholas Collins (School of Agriculture
528 Food and Wine, The University of Adelaide, Adelaide, SA, Australia) who supervised these projects and
529 thank all the technical staff involved in the field experiments. We are grateful for the insights offered in the
530 comments from anonymous reviewers and editors.

531

532 **Financial support.** The data used for this paper were derived from research projects funded by the Grains
533 Research and Development Corporation (GRDC) under project UA00123 and UA00147.

534

535 **Conflict of interest.** The authors declare there are no conflicts of interest.

536

537 **Ethical standard.** Not applicable.

538
539 **Author contributions.** Muhuddin Rajin Anwar (MRA), Livinus Emebiri (LE), Ryan H. L. Ip (RHLL),
540 David J. Lockett (DJL), Yashvir S. Chauhan (YSC) and Ketema T. Zeleke (KZ): Investigation;
541 Methodology; Data curation; Formal analysis; Writing – review & editing. MRA, DJL, RHLL:
542 conceptualised the model analysis and wrote the first draft of the manuscript, which all authors further
543 revised. LE supervised the field experimentations and data collection.

544

545 **References**

- 546 Abbate PE, Andrade FH and Culot JP (1995) The effects of radiation and nitrogen on number of grains
547 in wheat. *The Journal of Agricultural Science* **124**, 351-360.
548 <https://doi.org/10.1017/S0021859600073317>.
- 549 Abhinandan K, Skori L, Stanic M, Hickerson NMN, Jamshed M and Samuel MA (2018) Abiotic stress
550 signaling in wheat – an inclusive overview of hormonal interactions during abiotic stress
551 responses in wheat. *Frontiers in Plant Science* **9**, 734. <https://doi.org/10.3389/fpls.2018.00734>.
- 552 ABS (2024) Australian Bureau of Statistics 2021-22, Agricultural Commodities, Australia, ABS.
553 Available at <https://www.abs.gov.au/> (accessed 11 January 2024).
- 554 Akumaga U and Alderman PD (2019) Comparison of Penman–Monteith and Priestley-Taylor
555 Evapotranspiration Methods for Crop Modeling in Oklahoma. *Agronomy Journal* **111**, 1171-
556 1180. <https://doi.org/10.2134/agronj2018.10.0694>.
- 557 Anwar MR, Liu DL, Farquharson R, Macadam I, Abadi A, Finlayson J, Wang B and Ramilan T (2015)
558 Climate change impacts on phenology and yields of five broadacre crops at four climatologically
559 distinct locations in Australia. *Agricultural Systems* **132**, 133–144.
560 <https://doi.org/10.1016/j.agsy.2014.09.010>.
- 561 Anwar MR, Lockett DJ, Chauhan YS, Ip RHL, Maphosa L, Simpson M, Warren A, Raman R, Richards
562 MF, Pengilley G, Hobson K and Graham N (2022) Modelling the effects of cold temperature
563 during the reproductive stage on the yield of chickpea (*Cicer arietinum* L.). *International*
564 *Journal of Biometeorology* **66**, 111–125. <https://doi.org/10.1007/s00484-021-02197-8>.

- 565 APSIM (2023) SoilWat. Available at <https://www.apsim.info/documentation/model-documentation/soil->
566 [modules-documentation/soilwat/](https://www.apsim.info/documentation/model-documentation/soil-modules-documentation/soilwat/). (accessed 13 October 2023).
- 567 Asseng S, Ewert F, Martre P, Rötter RP, Lobell DB, Cammarano D, Kimball BA, Ottman MJ, Wall,
568 GW, White JW, Reynolds MP, Alderman PD, Prasad PVV, Aggarwal PK, Anothai J, Basso B,
569 Biernath C, Challinor AJ, De Sanctis G, Doltra J, Fereres E, Garcia-Vila M, Gayler S,
570 Hoogenboom G, Hunt LA, Izaurralde RC, Jabloun M, Jones CD, Kersebaum KC, Koehler A-K,
571 Müller C, Kumar SN, Nendel C, O’Leary G, Olesen JE, Palosuo T, Priesack E, Rezaei EE,
572 Ruane AC, Semenov MA, Shcherbak I, Stöckle C, Stratonovitch P, Streck T, Supit I, Tao F,
573 Thorburn PJ, Waha K, Wang E, Wallach D, Wolf J, Zhao Z and Zhu Y (2015) Rising
574 temperatures reduce global wheat production. *Nature Climate Change* **5**, 143–147.
575 <https://doi.org/10.1038/nclimate2470>.
- 576 Chapman SC, Hammer GL and Meinke H (1993) A sunflower simulation model. I. Model development.
577 *Agronomy Journal* **85**, 725–735. <https://doi.org/10.2134/agronj1993.00021962008500030038x>.
- 578 Chenu K, Cooper M, Hammer GL, Mathews KL, Dreccer MF and Chapman SC (2011) Environment
579 characterization as an aid to wheat improvement: interpreting genotype–environment
580 interactions by modelling water-deficit patterns in North-Eastern Australia. *Journal of*
581 *Experimental Botany* **62**, 1743–1755. <https://doi.org/10.1093/jxb/erq459>.
- 582 Collins N, Hildebrand S, Taylor K, Taylor H, Plemm D, Lohraseb I, Shirdelmoghanloo H, Erena M,
583 Rahman M, Taylor J, Munoz-Santa S, Mather D, Heuer S, Sissons M and Emebiri L (2017)
584 Understanding heat impacts on wheat to breed future tolerance. GRDC Update. Available at
585 [https://grdc.com.au/resources-and-publications/grdc-update-papers/tab-content/grdc-update-](https://grdc.com.au/resources-and-publications/grdc-update-papers/tab-content/grdc-update-papers/2017/02/understanding-heat-impacts-on-wheat-to-breed-future-tolerance)
586 [papers/2017/02/understanding-heat-impacts-on-wheat-to-breed-future-tolerance](https://grdc.com.au/resources-and-publications/grdc-update-papers/tab-content/grdc-update-papers/2017/02/understanding-heat-impacts-on-wheat-to-breed-future-tolerance). (accessed 22
587 October 2023).
- 588 Collins B and Chenu K (2021) Improving productivity of Australian wheat by adapting sowing date and
589 genotype phenology to future climate. *Climate Risk Management* **32**, 100300.
590 <https://doi.org/10.1016/j.crm.2021.100300>.
- 591 Didari S, Talebnejad R, Bahrami M and Mahmoudi MR (2023) Dryland farming wheat yield prediction
592 using the Lasso regression model and meteorological variables in dry and semi-dry region.
593 *Stochastic Environmental Research and Risk Assessment* **37**, 1-19.
594 <https://doi.org/10.1007/s00477-023-02490-5>.

- 595 Dormann CF, Elith J, Bacher S, Buchmann C, Carl G, Carré G, Marquéz JRG, Gruber B, Lafourcade B,
596 Leitão PJ, Münkemüller T, McClean C, Osborne PE, Reineking B, Schröder B, Skidmore AK,
597 Zurell D and Lautenbach S (2013) Collinearity: a review of methods to deal with it and a
598 simulation study evaluating their performance. *Ecography* **36**, 27-46.
599 <https://doi.org/10.1111/j.1600-0587.2012.07348.x>.
- 600 Emebiri L, Erena MF, Taylor K, Hildebrand S, Maccaferri M and Collins NC (2024) Field screening for
601 heat-stress tolerance of floret fertility in wheat (*Triticum aestivum* and *T. durum*). *Crop &*
602 *Pasture Science* **75**, CP23214. <https://doi.org/10.1071/CP23214>.
- 603 Erena MF, Lohraseb I, Munoz-Santa I, Taylor JD, Emebiri LC and Collins NC (2021) The WtmsDW
604 locus on wheat chromosome 2B controls major natural variation for floret sterility responses to
605 heat stress at booting stage. *Frontiers in Plant Science* **12**, 635397.
606 <https://doi.org/10.3389/fpls.2021.635397>.
- 607 FAO (2016) Food and Agriculture Organization of the United Nations. Climate change and food
608 security: Risks and responses. Available at <https://www.fao.org/3/i5188e/I5188E.pdf> (accessed 5
609 January 2024).
- 610 Fischer RA (1985) Number of kernels in wheat crops and the influence of solar radiation and
611 temperature. *The Journal of Agricultural Science* **105**, 447–461.
612 <https://doi.org/10.1017/S0021859600056495>.
- 613 Friedman JH, Hastie T and Tibshirani R (2010) Regularization Paths for Generalized Linear Models via
614 Coordinate Descent. *Journal of Statistical Software* **33**, 1–22.
615 <https://doi.org/10.18637/jss.v033.i01>.
- 616 GRDC GrowNote (2017) Grains Research and Development Corporation (GRDC), Durum Southern
617 Region. Available at [https://grdc.com.au/resources-and-publications/grownotes/crop-](https://grdc.com.au/resources-and-publications/grownotes/crop-agronomy/durum-southern-region-grownotes)
618 [agronomy/durum-southern-region-grownotes](https://grdc.com.au/resources-and-publications/grownotes/crop-agronomy/durum-southern-region-grownotes) and
619 [https://grdc.com.au/](https://grdc.com.au/data/assets/pdf_file/0016/301651/GRDC-GrowNotes-Durum-SOUTHERN.pdf)
620 [_data/assets/pdf_file/0016/301651/GRDC-GrowNotes-Durum-](https://grdc.com.au/data/assets/pdf_file/0016/301651/GRDC-GrowNotes-Durum-SOUTHERN.pdf)
[SOUTHERN.pdf](https://grdc.com.au/data/assets/pdf_file/0016/301651/GRDC-GrowNotes-Durum-SOUTHERN.pdf). (accessed 11 January 2024).
- 621 He D and Wang E (2019) On the relation between soil water holding capacity and dryland crop
622 productivity. *Geoderma* **353**, 11–24. <https://doi.org/10.1016/j.geoderma.2019.06.022>.

- 623 Hochman Z, Gobbett DL and Horan H (2017) Climate trends account for stalled wheat yields in
624 Australia since 1990. *Global Change Biology* **23**, 2071–2081. <https://doi.org/10.1111/gcb.13604>.
- 625 Holzworth DP, Huth NI, deVoil PG, Zurcher EJ, Herrmann NI, McLean G, Chenu K, van Oosterom EJ,
626 Snow V, Murphy C, Moore AD, Brown H, Whish JPM, Verrall S, Fainges J, Bell LW, Peake
627 AS, Poulton PL, Hochman Z, Thorburn PJ, Gaydon DS, Dalgliesh NP, Rodriguez D, Cox H,
628 Chapman S, Doherty A, Teixeira E, Sharp J, Cichota R, Vogeler I, Li FY, Wang E, Hammer GL,
629 Robertson MJ, Dimes JP, Whitbread AM, Hunt J, van Rees H, McClelland T, Carberry PS,
630 Hargreaves JNG, MacLeod N, McDonald C, Harsdorf J, Wedgwood S and Keating BA (2014)
631 APSIM - Evolution towards a new generation of agricultural systems simulation. *Environmental*
632 *Modelling & Software* **62**, 327–350. <http://dx.doi.org/10.1016/j.envsoft.2014.07.009>.
- 633 Isbell RF, National Committee on Soil and Terrain (2021) The Australian Soil Classification, third ed.
634 CSIRO Publishing, Melbourne.
- 635 Kerr RB, Hasegawa T, Lasco R, Bhatt I, Deryng D, Farrell A, Gurney-Smith H, Ju H, Lluch-Cota S,
636 Meza F, Nelson G, Neufeldt H, and Thornton P (2022) Food, Fibre, and Other Ecosystem
637 Products. In Pörtner HO, Roberts DC, Tignor M, Poloczanska ES, Mintenbeck K, Alegría A,
638 Craig M, Langsdorf S, Löschke S, Möller V, Okem A and Rama B (eds). *Climate Change 2022:*
639 *Impacts, Adaptation and Vulnerability. Contribution of Working Group II to the Sixth*
640 *Assessment Report of the Intergovernmental Panel on Climate Change*, pp. 713–906. Cambridge
641 University Press, Cambridge, UK and New York, NY, USA. Available at
642 <https://doi.org/10.1017/9781009325844.007>.
- 643 Kutner MH, Nachtsheim CJ, Neter J and Li W (2005) *Applied linear statistical models* (Vol. 5). Boston:
644 McGraw-Hill Irwin.
- 645 Li S, Wang B, Feng P, Liu DL, Li C, Shi L and Yu Q (2022) Assessing climate vulnerability of
646 historical wheat yield in south-eastern Australia’s wheat belt. *Agricultural Systems* **196**, 103340.
647 <https://doi.org/10.1016/j.agsy.2021.103340>.
- 648 Liu DL, Anwar MR, O’Leary G and Conyers MK (2014) Managing wheat stubble as an effective
649 approach to sequester soil carbon in a semi-arid environment: Spatial modelling. *Geoderma*
650 **214–215**, 50–61. <http://dx.doi.org/10.1016/j.geoderma.2013.10.003>.
- 651 Lobell DB, Schlenker W and Costa-Roberts J (2011) Climate Trends and Global Crop Production Since
652 1980. *Science* **333**, 616–620. <https://doi.org/10.1126/science.1204531>.

- 653 Mittler R (2006) Abiotic stress, the field environment and stress combination. *Trends in Plant Science*
654 **11**, 15–19; <https://doi.org/10.1016/j.tplants.2005.11.002>.
- 655 Monahan JF (2011) A Primer on Linear Models. Chapman and Hall/CRC, New York.
656 <https://doi.org/10.1201/b11551>.
- 657 Nuttall J, Brady S, Brand J, O’Leary G and Fitzgerald GJ (2012) Heat waves and wheat growth under a
658 future climate. In Yunusa I (ed). Capturing Opportunities and Overcoming Obstacles in
659 Australian Agronomy. Proceedings of 16th Australian Agronomy Conference 2012, 14-18
660 October 2012, Armidale, NSW. Available at [http://www.regional.org.au/au/asa/2012/climate-](http://www.regional.org.au/au/asa/2012/climate-change/8085_nuttalljg.htm)
661 [change/8085_nuttalljg.htm](http://www.regional.org.au/au/asa/2012/climate-change/8085_nuttalljg.htm).
- 662 Nuttall JG, Barlow KM, Delahunty AJ, Christy BP and O’Leary GJ (2018) Acute High Temperature
663 Response in Wheat. *Agronomy Journal* **110**, 1296-1308.
664 <https://doi.org/10.2134/agronj2017.07.0392>.
- 665 OEC (2023) The Observatory of Economic Complexity. Australia (AUS) Exports, Imports, and Trade
666 Partners. Available at <https://oec.world/en/profile/country/aus?yearlyTradeFlowSelector=flow1>.
667 (accessed 11 January 2024).
- 668 OECD/FAO (2023) OECD-FAO Agricultural Outlook. Paris: OECD Agriculture Statistics. Available at
669 <https://www.oecd.org/publications/oecd-fao-agricultural-outlook-19991142.htm>. (accessed 12
670 October 2023).
- 671 Ortiz-Monasterio JI, Dhillon SS and Fischer RA (1994) Date of sowing effects on grain-yield and yield
672 components of irrigated spring wheat cultivars and relationships with radiation and temperature
673 in Ludhiana, India. *Field Crops Research* **37**, 169–184. [https://doi.org/10.1016/0378-](https://doi.org/10.1016/0378-4290(94)90096-5)
674 [4290\(94\)90096-5](https://doi.org/10.1016/0378-4290(94)90096-5).
- 675 Pandey P, Irulappan V, Bagavathiannan MV and Senthil-Kumar M (2017) Impact of Combined Abiotic
676 and Biotic Stresses on Plant Growth and Avenues for Crop Improvement by Exploiting Physio-
677 morphological Traits. *Frontiers in Plant Science* **8**, 537. <https://doi.org/10.3389/fpls.2017.00537>.
- 678 Paredes P, Rodrigues GC, Alves I and Pereira LS (2014) Partitioning evapotranspiration, yield
679 prediction, and economic returns of maize under various irrigation management strategies.
680 *Agricultural Water Management* **135**, 27–39. <https://doi.org/10.1016/j.agwat.2013.12.010>.

- 681 Prasad PVV, Pisipati SR, Momcilovic I and Ristic Z (2011) Independent and combined effects of high
682 temperature and drought stress during grain filling on plant yield and chloroplast EF-Tu
683 expression in spring wheat. *Journal of Agronomy and Crop Science* **197**, 430–441;
684 <https://doi.org/10.1111/j.1439-037X.2011.00477.x>.
- 685 R Core Team (2023) R: A Language and Environment for Statistical Computing, Vienna, Austria.
686 Available at <http://www.R-project.org>.
- 687 Ru C, Hu X, Chen D, Wang W, Zhen J and Song T (2023) Individual and combined effects of heat and
688 drought and subsequent recovery on winter wheat (*Triticum aestivum* L.) photosynthesis,
689 nitrogen metabolism, cell osmoregulation, and yield formation. *Plant Physiology and*
690 *Biochemistry* **196**, 222-235. <https://doi.org/10.1016/j.plaphy.2023.01.038>.
- 691 Sinha R, Fritschi FB, Zandalinas SI and Mittler R (2021) The impact of stress combination on
692 reproductive processes in crops. *Plant Science* **311**, 111007.
693 <https://doi.org/10.1016/j.plantsci.2021.111007>.
- 694 RStudio Team (2023) RStudio: Integrated Development for R. RStudio, PBC, Boston, MA URL.
695 Available at <https://posit.co/downloads/>.
- 696 Sadras VO and McDonald G (2012) Water use efficiency of grain crops in Australia: principles,
697 benchmarks and management. GRDC Project Code: DAS00089. Available at
698 [https://grdc.com.au/__data/assets/pdf_file/0030/159186/grdcpublicationwateruseefficiencyofgrai](https://grdc.com.au/__data/assets/pdf_file/0030/159186/grdcpublicationwateruseefficiencyofgraincropsinaustraliapdf.pdf)
699 [ncropsinaustraliapdf.pdf](https://grdc.com.au/__data/assets/pdf_file/0030/159186/grdcpublicationwateruseefficiencyofgraincropsinaustraliapdf.pdf).
- 700 Shafiee S, Lied LM, Burud I, Dieseth JA, Alsheikh M and Lillemo M (2021) Sequential forward
701 selection and support vector regression in comparison to LASSO regression for spring wheat
702 yield prediction based on UAV imagery. *Computers and Electronics in Agriculture* **183**, 106036.
703 <https://doi.org/10.1016/j.compag.2021.106036>.
- 704 Shirdelmoghanloo H, Taylor JD, Lohraseb I, Rabie H, Brien C, Timmins A, Martin P, Mather DE,
705 Emebiri L and Collins NC (2016) A QTL on the short arm of wheat (*Triticum aestivum* L.)
706 chromosome 3B affects the stability of grain weight in plants exposed to a brief heat shock early
707 in grain filling. *BMC Plant Biology* **16**, 100. <https://doi.org/10.1186/s12870-016-0784-6>.

- 708 Sissons M, Fleming D, Taylor JD, Emebiri L, Eckermann P, Collins NC (2024; *submitted*). Effects of
709 heat exposure from late sowing on agronomic traits and the technological quality of hexaploid
710 wheat. *Journal of Cereal Science* (Submitted).
- 711 Sissons M, Fleming D, Taylor JD, Emebiri L and Collins NC (2018) Effects of heat exposure from late
712 sowing on the agronomic and technological quality of tetraploid wheat. *Cereal Chemistry* **95**,
713 274-287. <https://doi.org/10.1002/cche.10027>.
- 714 Slafer GA, Savin R and Sadras VO (2023) Wheat yield is not causally related to the duration of the
715 growing season. *European Journal of Agronomy* **148**, 126885.
716 <https://doi.org/10.1016/j.eja.2023.126885>.
- 717 Stone PJ and Nicolas ME (1994) Wheat cultivars vary widely in their responses of grain yield and
718 quality to short periods of post-anthesis heat stress. *Australian Journal of Plant Physiology* **21**,
719 887–900. <http://dx.doi.org/10.1071/PP9940887>.
- 720 Street K, Bari A, Mackay M and Amri A (2016) How the Focused Identification of Germplasm Strategy
721 (FIGS) is used to mine plant genetic resources collections for adaptive traits. In Maxted N,
722 Dulloo ME, Ford-Lloyd BV (Eds.). *Enhancing crop genepool use: capturing wild relative and*
723 *landrace diversity for crop improvement*. CAB International, Wallingford, UK, pp. 54–63.
- 724 Talukder ASMHM, McDonald GK and Gill GS (2013) Effect of short-term heat stress prior to flowering
725 and at early grain set on the utilization of water-soluble carbohydrate by wheat genotypes. *Field*
726 *Crops Research*, **147**, 1–11. <https://doi.org/10.1016/j.fcr.2013.03.013>.
- 727 Talukder SK, Babar MA, Vijayalakshmi K, Poland J, Prasad VV, Bowden R and Fritz A (2014)
728 Mapping QTL for the traits associated with heat tolerance in wheat (*Triticum aestivum* L.). *BMC*
729 *Genetics* **15**, 97. <https://doi.org/10.1186/s12863-014-0097-4>.
- 730 Teixeira EI, Fischer G, van Velthuizen H, Walter C and Ewert F (2013) Global hot-spots of heat stress
731 on agricultural crops due to climate change. *Agricultural and Forest Meteorology* **170**, 206–215.
732 <https://doi.org/10.1016/j.agrformet.2011.09.002>.
- 733 Tibshirani R (1996) Regression Shrinkage and Selection via the Lasso. *Journal of the Royal Statistical*
734 *Society: Series B (methodological)* **58**, 267–288. [https://doi.org/10.1111/j.2517-](https://doi.org/10.1111/j.2517-6161.1996.tb02080.x)
735 [6161.1996.tb02080.x](https://doi.org/10.1111/j.2517-6161.1996.tb02080.x).

- 736 Trethowan RM (2022) Abiotic Stresses. In Reynolds MP and Braun HJ (eds). Wheat Improvement.
737 Springer, Cham. https://doi.org/10.1007/978-3-030-90673-3_10.
- 738 Trout TJ and DeJonge KC (2017) Water productivity of maize in the US high plains. *Irrigation Science*
739 **35**, 251–266. <https://doi.org/10.1007/s00271-017-0540-1>.
- 740 Unkovich M, Baldock J and Farquharson, R (2018) Field measurements of bare soil evaporation and
741 crop transpiration, and transpiration efficiency, for rainfed grain crops in Australia – A review.
742 *Agricultural Water Management* **205**, 72–80. <https://doi.org/10.1016/j.agwat.2018.04.016>.
- 743 Unkovich M, McBeath T, Moodie M and Macdonald LM (2023) High soil strength and cereal crop
744 responses to deeper tillage on sandy soils in a semi-arid environment. *Field Crops Research* **291**,
745 108792. <https://doi.org/10.1016/j.fcr.2022.108792>.
- 746 Wang B, Liu DL, Asseng S, Macadam I and Yu Q (2017) Modelling wheat yield change under CO₂
747 increase, heat and water stress in relation to plant available water capacity in eastern Australia.
748 *European Journal of Agronomy* **90**, 152-161. <https://doi.org/10.1016/j.eja.2017.08.005>.
- 749 Wickham H, Averick M, Bryan J, Chang W, McGowan LD, François R, Grolemund G, Hayes A, Henry
750 L, Hester J, Kuhn M, Pedersen TL, Miller E, Bache SM, Müller K, Ooms J, Robinson D, Seidel
751 DP, Spinu V, Takahashi K, Vaughan D, Wilke C, Woo K and Yutani H (2019) Welcome to the
752 tidyverse. *Journal of Open Source Software* **43**, 1686. <https://doi.org/10.21105/joss.01686>.
- 753 Xing H, Liu DL, Li G, Wang B, Anwar MR, Crean J, Lines-Kelly R and Yu G (2017) Incorporating
754 grain legumes in cereal-based cropping systems to improve profitability in southern New South
755 Wales, Australia. *Agricultural Systems* **154**, 112–123.
756 <https://doi.org/10.1016/j.agsy.2017.03.010>.
- 757 Yang Y, Liu DL, Anwar MR, O’Leary G, Macadam I and Yang Y (2016). Water use efficiency and crop
758 water balance of rainfed wheat in a semi-arid environment: sensitivity of future changes to
759 projected climate changes and soil type. *Theoretical and Applied Climatology* **123**, 565–579.
760 <https://doi.org/10.1007/s00704-015-1376-3>.
- 761 Zeleke K and Nendel C (2019) Growth and yield response of faba bean to soil moisture regimes and
762 sowing dates: Field experiment and modelling study. *Agricultural Water Management* **213**,
763 1063-1077. <https://doi.org/10.1016/j.agwat.2018.12.023>.

764 Zeleke KT, Anwar MR, Emebiri L and Lockett D (2023) Weather indices during reproductive phase
765 explain wheat yield variability. *The Journal of Agricultural Science* **161**, 617-632.
766 <https://doi.org/10.1017/S0021859623000503>.

767 Zhao C, Liu B, Piao S, Wang X, Lobell DB, Huang Y, Huang M, Yao Y, Bassu S, Ciais P, Durand J-L,
768 Elliott J, Ewert F, Janssens IA, Li T, Lin E, Liu Q, Martre P, Müller C, Peng S, Penuelas J,
769 Ruane AC, Wallach D, Wang T, Wu D, Liu Z, Zhu Y, Zhu Z and Asseng S (2017). Temperature
770 increase reduces global yields of major crops in four independent estimates. *Proceedings of the*
771 *National Academy of Sciences of the United States of America* **114**, 35.
772 <https://doi.org/10.1073/pnas.1701762114>.

773

774

Tables and Figures

Table 1. Characterisation of the soils, site descriptions, and long term (1950-2022) average climate variables for the two sites used in the field experiments described in this paper. PAWC= plant available water capacity; BD=bulk density; OC= soil organic carbon; GS= growing season (April to October); maxT= mean annual maximum temperature; minT= mean annual minimum temperature; avT=mean annual average temperature; Frost= mean annual any day where minT \leq 0°C. * = coefficient of variation (CV, %) in parentheses. ^s = top soil layer equals 0-10 cm.

Site	Latitude Longitude	Soil type	Soil profile		Top soil layer ^s			rain (mm)	GS rain (mm)	Solar (MJ/m ²)	GS Solar (MJ/m ²)	maxT GS (°C)	MinT GS (°C)	avT GS (°C)	Frost GS (days)
			Soil depth (m)	Total PAWC (mm)	BD (g/cc)	pH (1:5 water)	OC (%)								
Leeton	34.73 S	Wunnamurra Clay (clay loam)	1.80	293	1.30	7.20	1.75	478	293	17.7	13.2	18.5	6.3	12.4	22
	146.55 E							(32.5)*	(37.3)	(5.25)	(6.78)	(4.39)	(9.57)	(3.71)	(44.4)
Wagga Wagga	35.05 S	Kandosol	1.25	128	1.45	6.43	1.69	560	339	17.3	12.8	17.4	5.7	11.6	28
	147.35 E							(29.4)	(34.6)	(5.25)	(6.79)	(5.17)	(11.89)	(4.02)	(42.6)

Table 2: Mean grain yield and grain weight of six wheat germplasm groupings (across two wheat species, bread wheat and durum) sown at one or two different times. The number of genotypes of each germplasm category and the range of genotype means for grain yield (t/ha) and 1000-grain weight (TGW, g) are also presented. na = not available.

Category	Sowing time	Number of genotypes	Mean sowing date (Julian day)	Mean yield (t/ha)	Range in yield (t/ha)	Mean TGW (g)	Range in TGW (g)
BreadWheat_ABDLines	1	72	156	5.66	3.17 - 8.96	41.9	26.80 - 50.38
BreadWheat_ABDLines	2	61	217	3.85	1.86 - 5.75	33.3	21.38 - 42.65
BreadWheat_Elite	1	217	156	5.33	2.13 - 8.73	39.1	27.92 - 52.25
BreadWheat_Elite	2	219	217	3.33	0.19 - 5.46	31.2	na
BreadWheat_Landraces	1	196	157	3.74	2.00 - 6.35	40.3	30.68 - 58.0
BreadWheat_Landraces	2	201	218	2.07	0.44 - 4.23	31.9	22.28 - 48.40
BreadWheat_NILines	1	61	137	2.62	2.30 - 2.92	33.6	25.52 - 39.30
Durum_Biparent	1	232	152	2.78	2.25 - 3.43	35.1	26.51 - 44.20
Durum_Biparent	2	322	216	1.48	0.88 - 2.03	40.4	31.82 - 47.76
Durum_Elite	1	4	152	4.73	2.59 - 7.22	42.0	38.34 - 45.30
Durum_Elite	2	6	215	2.49	0.86 - 4.61	39.8	32.05 - 44.95

Table 3. Water use and water-use efficiency based on transpiration (Tran and wue_Tran) and evapotranspiration (ET and wue_ET), respectively for six wheat germplasm groupings sown at one or two different times. Overall mean values are presented along with corresponding ranges for individual genotype means.

Category	Sowing time	Mean	Range in	Mean	Range in	Mean	Range in	Mean	Range in
		Tran	Tran	ET	ET	wue_Tran	wue_Tran	wue_ET	wue_ET
		(mm)				(Kg grain/ha/mm)			
BreadWheat_ABDLines	1	155	136 - 187	245	219 - 277	36.6	20.1 - 49.9	23.0	13.7 - 33.5
BreadWheat_ABDLines	2	222	194 - 255	308	272 - 346	17.3	7.8 - 23.1	12.5	5.8 - 16.8
BreadWheat_Elite	1	158	136 - 188	249	219 - 299	33.9	15.6 - 57.5	21.4	9.1 - 34.9
BreadWheat_Elite	2	222	187 - 258	310	262 - 363	15.2	0.92 - 24.6	10.9	0.6 - 17.6
BreadWheat_Landraces	1	157	136 - 188	260	219 - 313	24.5	13.7 - 46.6	14.6	7.9 - 28.2
BreadWheat_Landraces	2	220	191 - 258	313	268 - 370	9.7	1.69 - 21.7	6.8	1.2 - 15.2
BreadWheat_NILines	1	159	158 - 159	269	265 - 270	16.5	14.4 - 18.4	9.7	8.5 - 10.8
Durum_Biparent	1	146	138 - 156	257	239 - 287	18.9	15.8 - 21.9	10.8	9.1 - 12.1
Durum_Biparent	2	189	141 - 232	279	223 - 332	7.8	5.7 - 9.9	5.3	3.7 - 6.4
Durum_Elite	1	163	142 - 186	258	245 - 275	27.8	18.3 - 38.9	17.9	10.6 - 26.3
Durum_Elite	2	211	151 - 254	299	236 - 342	11.1	5.7 - 18.2	7.9	3.6 - 13.6

Table 4. Pearson correlation coefficients between yield, 1000 grain weight (TGW), evapotranspiration (ET), transpiration (Tran), accumulated water deficit (SDR = supply-demand ratio), and heat stress (H>30 = number of days with temperatures >30 °C) during growth Period 3 (see text for explanation) in the “BreadWheat_ABDLine” germplasm category for both sites (Wagga Wagga and Leeton) combined, including all sowing times. Significance levels are indicated as follows: * 0.01 < p < 0.05, ** 0.001 < p < 0.01, *** p < 0.001

BreadWheat_ABDLines germplasm group						
	Yield	TGW	ET	Tran	SDR	H>30
Yield		0.52***	-0.38***	-0.37***	-0.15**	-0.62***
TGW			-0.61***	-0.62***	0.16**	-0.60***
ET				0.99***	-0.59***	0.70***
Tran					-0.60***	0.68***
SDR						-0.23***

Table 5. Pearson correlation coefficients between yield, 1000 grain weight (TGW), evapotranspiration (ET), transpiration (Tran), accumulated water deficit (SDR = supply-demand ratio), and heat stress (H>30 = number of days with temperatures >30 °C) during growth Period 3 (see text for explanation) in the “BreadWheat_Elite” germplasm category for both sites combined, including all sowing times. Significance levels are indicated as follows: * 0.01 < p < 0.05, ** 0.001 < p < 0.01, *** p < 0.001

BreadWheat_Elite germplasm group						
	Yield	TGW	ET	Tran	SDR	H>30
Yield		0.59***	-0.62***	-0.66***	0.20***	-0.73***
TGW			-0.28***	-0.37***	-0.11*	-0.49***
ET				0.97***	-0.61***	0.82***
Tran					-0.55***	0.77***
SDR						-0.29***

Table 6. Pearson correlation coefficients between yield, 1000 grain weight (TGW), evapotranspiration (ET), transpiration (Tran), accumulated water deficit (SDR = supply-demand ratio), and heat stress (H>30 = number of days with temperatures >30 °C) during growth Period 3 (see text for explanation) in the “BreadWheat_Landraces” germplasm category for both sites combined, including all sowing times. Significance level are indicated as follows: * 0.01 < p < 0.05, ** 0.001 < p < 0.01, *** p < 0.001

BreadWheat_Landraces germplasm group						
	Yield	TGW	ET	Tran	SDR	H>30
Yield		-0.10	0.24	0.27	-0.93***	-0.85**
TGW			-0.59	-0.56	0.25	0.26
ET				0.99***	-0.33	-0.18
Tran					-0.32	-0.13

SDR

0.97***

Table 7. Pearson correlation coefficients between yield, 1000 grain weight (TGW), evapotranspiration (ET), transpiration (Tran), accumulated water deficit (SDR = supply-demand ratio), and heat stress (H>30 = number of days with temperatures >30 °C) during growth Period 3 (see text for explanation) in the “BreadWheat_NILines” germplasm category for both sites combined, including all sowing times. Significance level are indicated as follows: * 0.01 < p < 0.05, **0.001 < p < 0.01, *** p < 0.001

BreadWheat_NILines germplasm group						
	Yield	TGW	ET	Tran	SDR	H>30
Yield		0.57***	-0.18*	-0.20*	-0.40***	-0.30***
TGW			-0.31***	-0.34***	-0.34***	-0.07
ET				0.99***	-0.50***	0.36**
Tran					-0.48***	0.30***
SDR						-0.40***

Table 8. Pearson correlation coefficients between yield, 1000 grain weight (TGW), evapotranspiration (ET), transpiration (Tran), accumulated water deficit (SDR = supply-demand ratio), and heat stress (H>30 = number of days with temperatures >30 °C) during growth Period 3 (see text for explanation) in the “Durum_Biparent” germplasm category for both sites combined, including all sowing times. Significance level are indicated as follows: * 0.01 < p < 0.05, **0.001 < p < 0.01, *** p < 0.001

Durum_Biparent germplasm group						
	Yield	TGW	ET	Tran	SDR	H>30
Yield		0.16	-0.06	-0.06	0.13	
TGW			0.06	0.03	0.05	
ET				0.90***	0.33**	
Tran					0.34**	
SDR						

Table 9. Pearson correlation coefficients between yield, 1000 grain weight (TGW), evapotranspiration (ET), transpiration (Tran), accumulated water deficit (SDR = supply-demand ratio), and heat stress (H>30 = number of days with temperatures >30 °C) during growth Period 3 (see text for explanation) in the “Durum_Elite” germplasm category for both sites combined, including all sowing times. Significance level are indicated as follows: * 0.01 < p < 0.05, ** 0.001 < p < 0.01, *** p < 0.001

Durum_Elite germplasm group						
	Yield	TGW	ET	Tran	SDR	H>30
Yield	1.00	-0.75***	0.18***	-0.16***	-0.22***	-0.87***

TGW	1.00	-0.17***	0.11*	0.08	0.62***
ET		1.00	0.93***	-0.07	0.05
Tran			1.00	-0.10*	0.31***
SDR				1.00	0.57***

Table 10. Estimated coefficients from LASSO (least absolute shrinkage and selection operator) and model performance for Leeton and Wagga Wagga (all sowing times pooled), as measured by estimated coefficients, RMSE, Adjusted (Adj) R², and Lambda (λ). The explanatory variables consisted of water stress (D), heat stress (H; number of days with temperatures >30°C), evapotranspiration (ET; Priestley-Taylor method), and the response variable was wheat grain yield (t/ha) for each of the four growing periods (period 1 - 4, see text for details) *Explanatory variables (D, H, or ET) plus period number

BreadWheat_ABDLines		BreadWheat_Elite	
Item	LASSO coefficients	Item	LASSO coefficients
Intercept	4.834	Intercept	4.327
D_period1*	-0.498		-0.036
D_period2	-0.341		-0.501
D_period3	-0.789		0
D_period4	0		-0.261
H_period1	-0.575		-0.227
H_period2	-0.432		-0.449
H_period3	-0.073		-0.337
H_period4	-0.312		-0.227
ET_period1	0.431		0.344
ET_period2	0		0
ET_period3	0		0
ET_period4	0		0
	RMSE 0.846		0.976
	Adj R ² 0.66		0.57
	λ 0.0011		0.0054
BreadWheat_Landraces		Durum_Biparent	
Item	LASSO coefficients	Item	LASSO coefficients
Intercept	2.893	Intercept	2.027
D_period1*	-0.102		-0.207

D_period2	0	-0.241
D_period3	-0.064	0
D_period4	0	0
H_period1	-0.143	0
H_period2	-0.725	-0.361
H_period3	0	-0.215
H_period4	-0.247	-0.135
ET_period1	0.161	0.224
ET_period2	0	0
ET_period3	0	0
ET_period4	0	0.278
RMSE	0.605	0.110
Adj R ²	0.70	0.98
λ	0.0063	0.000

Figures:




Figure 1. Boxplots of mean SDR (supply-demand ratio = water stress) for individual genotypes within six germplasm groups and for each of the growing periods (see text for details). Data includes all sowing times. The number of genotypes in each group and sowing time is given in Table 3.




Figure 2. Boxplots of potential evapotranspiration for individual genotypes within six germplasm groups and for each of the growing periods (see text for details). Data includes all sowing times. The number of genotypes in each group and sowing time is given in Table 3.




Figure 3. Boxplots of heat stress for individual genotypes within six germplasm groups and for each of the growing periods (see text for details). Data includes all sowing times. The number of genotypes in each group and sowing time is given in Table 3.




Figure 4. Plots of predicted wheat yield (t/ha) derived from LASSO modelling against observed yield for four wheat germplasm categories. The best fitted straight line, together with the R^2 and P-value of the slope term, are provided for better visual assessment.

Table S2. Details (means and ranges) of the “sowing-date to flowering-date interval” (in days) for the genotypes within each germplasm category and across all combinations of Site, Year, and Sowing-time (1 = ‘early’, 2 = ‘late’).

Germplasm category	Site	Year	Sowing time	Number of genotypes	Mean sowing to flowering interval (days)	Range in sowing to flowering interval (days)
BreadWheat_ABDLines	Leeton	2011	1	28	114.8	110.5 – 120.0
BreadWheat_ABDLines	Leeton	2011	2	28	84.2	79.3 – 86.7
BreadWheat_ABDLines	Leeton	2015	1	11	129.6	127.7 – 130.6
BreadWheat_ABDLines	Wagga Wagga	2012	1	33	122.7	117.3 – 133.4
BreadWheat_ABDLines	Wagga Wagga	2012	2	33	84.5	79.7 – 90.7
BreadWheat_Elite	Leeton	2011	1	98	116.7	110.1 – 136.4
BreadWheat_Elite	Leeton	2011	2	97	85.7	77.7 – 98.0
BreadWheat_Elite	Leeton	2015	1	6	130.2	128.5 – 132.0
BreadWheat_Elite	Leeton	2015	2	8	104.4	101.7 – 106.1
BreadWheat_Elite	Wagga Wagga	2012	1	111	124.2	116.9 – 137.5
BreadWheat_Elite	Wagga Wagga	2012	2	114	85.5	72.9 – 95.7
BreadWheat_Elite	Wagga Wagga	2018	1	2	138.6	137.1 – 140.1
BreadWheat_Landraces	Leeton	2011	1	77	133.0	112.6 – 148.2
BreadWheat_Landraces	Leeton	2011	2	80	94.2	83.3 – 107.9

Table S2. Details (means and ranges) of the “sowing-date to flowering-date interval” (in days) for the genotypes within each germplasm category and across all combinations of Site, Year, and Sowing-time (1 = ‘early’, 2 = ‘late’).

Germplasm category	Site	Year	Sowing time	Number of genotypes	Mean sowing to flowering interval (days)	Range in sowing to flowering interval (days)
BreadWheat_Landraces	Wagga Wagga	2012	1	119	134.0	115.5 – 148.1
BreadWheat_Landraces	Wagga Wagga	2012	2	121	91.1	76.0 – 107.8
BreadWheat_NILines	Wagga Wagga	2018	1	61	139.5	137.7 – 140.5
Durum_Biparent	Wagga Wagga	2018	1	71	138.1	137.3 – 138.9
Durum_Biparent	Wagga Wagga	2018	2	161	90.2	87.9 – 94.4
Durum_Biparent	Wagga Wagga	2019	1	161	115.6	109.6 – 120.3
Durum_Biparent	Wagga Wagga	2019	2	161	77.3	72.4 – 82.5
Durum_Elite	Leeton	2011	1	2	115.6	113.4 – 117.8
Durum_Elite	Leeton	2011	2	2	83.0	81.3 – 84.7
Durum_Elite	Wagga Wagga	2018	2	2	89.8	88.7 – 90.8
Durum_Elite	Wagga Wagga	2019	1	2	116.4	116.3 – 116.4
Durum_Elite	Wagga Wagga	2019	2	2	77.7	77.6 – 77.7

