

Calling time on the imperial lawn and the imperative for greenhouse gas mitigation

Len N. Gillman¹ , Barbara Bollard¹ and Sebastian Leuzinger² 

¹Drone Lab, Auckland University of Technology, Auckland, New Zealand and ²Department of Environmental Science, Auckland University of Technology, Auckland, New Zealand

Intelligence Briefing

Cite this article: Gillman LN, Bollard B, Leuzinger S (2023). Calling time on the imperial lawn and the imperative for greenhouse gas mitigation. *Global Sustainability* 6, e3, 1–9. <https://doi.org/10.1017/sus.2023.1>

Received: 2 June 2022
Revised: 22 December 2022
Accepted: 28 December 2022

Keywords:

above ground biomass; carbon sequestration; grass; greenhouse gas emissions; lawn; soil carbon; soil; tree; turf

Author for correspondence:

Len N. Gillman, E-mail: len.gillman@aut.ac.nz

Non-technical summary. As green spaces, lawns are often thought to capture carbon from the atmosphere. However, once mowing, fertilising and irrigation are taken into account, we show that they become carbon sources, at least in the long run. Converting unused urban and rural lawn and grassland to trees can make a substantial contribution to reducing greenhouse gas emissions and increasing carbon absorption from the atmosphere. However, it is imperative for governing bodies to put in place appropriate policies and incentives in order to achieve this.

Technical summary. Mown grass or lawn is a ubiquitous form of vegetation in human-dominated landscapes and it is often claimed to perform an ecosystem service by sequestering soil carbon. If lawn maintenance is included, however, we show that lawns become net carbon emitters. We estimate that globally, if one-third of mown grass in cities was returned to trees, 310–1630 million tonnes of carbon could be absorbed from the atmosphere, and up to 43 tonnes of carbon equivalent per hectare of emissions could be avoided over a two-decade time span. We therefore propose that local and central governments introduce policies to incentivise and/or regulate the conversion of underutilised grass into trees.

Social media summary. If unused lawns were planted with trees, a gigaton of carbon could be removed from the atmosphere over two decades.

1. Introduction

Globally, mown grass is one of the most common features of our human-shaped landscapes; it is almost ubiquitous in suburban residential gardens, especially in the USA, Australia and New Zealand (Ignatieva & Stewart, 2009). It dominates in parks (e.g. 75–95% of the park area in Europe (Gilbert, 1989)), covers approximately 23% of urban areas (Ignatieva & Hedblom, 2018) and is common on highway margins and street verges. Mown grass is estimated to cover 16.4 Mha in the contiguous 48 states of the USA alone (Milesi et al., 2005), an area which roughly equals that of England and Belgium combined. Globally it occupies 15–80 Mha of land within urban areas (Ignatieva & Hedblom, 2018). The imperial lawn originated from hand-cut grass in lieu of grazed pasture to demonstrate wealth among the gentry. It then became part of the order that colonists implanted on the conquered land, representing a pastoral nostalgia of the European landscape (Ignatieva & Stewart, 2009). Today a large range of lawns are formed from mown grass. While some of these can have a specific purpose (such as for sport fields and recreation), often their occurrence is purely historical because land that has no specific designated use is often mown just to keep it tidy, and in many cases, there is a lack of rationale for maintaining them as mown grass.

Mown grass has been promoted as providing important ecosystem services such as carbon sequestration, and while it may have advantages relative to impervious surfaces (Ignatieva & Hedblom, 2018; Velasco et al., 2021), we believe the more apposite comparison is with trees and shrublands that would have once occupied current lawns. Given the widespread nature of mown grass, and with humanity facing a catastrophic climate crisis, it is important to review the role lawns play with respect to greenhouse gases. Here we review the potential of carbon storage in mown grass relative to that provided by other vegetation types. Carbon sequestration/emissions and long-term storage estimates are projected over two decades, the time frame that aligns with the critical period for action on climate change. We conclude that mown grass in almost all cases makes a negative contribution (carbon release) when emissions associated with lawn maintenance are considered. Other forms of vegetation cover, such as shrubs and trees, store substantially more carbon, both below and above ground. We therefore recommend radical changes in policy settings to maximise conversion of mown grass to shrub, tree or mixed vegetation.

2. Methods

Articles reporting carbon sequestration in mown grass and planted trees were searched using Google scholar using the following search strings: (1) 'carbon sequestration' AND (grass OR

© The Author(s), 2023. Published by Cambridge University Press. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.



CrossMark

<https://doi.org/10.1017/sus.2023.1> Published online by Cambridge University Press

turf OR lawn) (2) 'carbon sequestration' AND tree*. Article titles and abstracts were examined for reporting of quantified carbon sequestration rates or storage. Studies were excluded if they only reported one component of carbon flux or if they related to sports fields or golf courses. References in selected publications were also examined for relevant publications. Summary data from 65 studies are reported in Supplementary Tables S1 and S2.

3. Carbon storage and emissions in lawns

Soil organic carbon (SOC) capacity is influenced by bioclimatic conditions and increases at higher latitudes due to slow mineralisation at low temperatures (Vasenev & Kuzyakov, 2018). Across the USA, SOC stored in the top 15 cm of residential lawns averaged 45.8 t ha^{-1} (Selhorst & Lal, 2013) (Table 1). In the top 30 cm of fertilised and irrigated lawn soil at a warm temperate site as much as 108 t ha^{-1} has been recorded (Weissert *et al.*, 2016). Reported annual gross carbon sequestration into lawn soils, summed over two decades, may be as high as 28 t ha^{-1} (Table 1). However, although sequestration of carbon into lawn soils can occur over some years, it can be expected to asymptote to zero within 30–50 years, depending on climate (Lindén *et al.*, 2020; Qian *et al.*, 2003). At very slow rates of accumulation, this might extend to 100 years (Smith *et al.*, 2018), after which carbon sequestration will level off. On the other hand, climate forcing gas emissions that occur through mowing, fertilising and irrigation are summative, and extend past periods of biological carbon sequestration into soils. Therefore, these need to be subtracted from gross sequestration to establish net carbon equivalent fluxes and pools.

The rate of carbon emissions due to mowing depends on the size and type of mower used and the regularity of mowing. Estimates range from 1.4 to 6.7 t C ha^{-1} over two decades (Table 1). Given that the estimated area of mown grass in the continental USA is 16.4 million hectares (Milesi *et al.*, 2005), this implies that in the USA alone 1.1–5.5 million tonnes of carbon are emitted every year due to mowing.

The addition of fertiliser to lawns causes emissions of N_2O , which has a climate forcing effect 298 times that of CO_2 (Townsend-Small & Czimczik, 2010a). Emissions from fertilising depend on the regularity and rate of application, but with the high rates often recommended, they can offset carbon sequestration over two decades by up to 26.6 tonnes carbon equivalent ha^{-1} , an amount almost equal to the highest rates of sequestration reported (Townsend-Small & Czimczik, 2010b) (Table 1).

Irrigation of lawns involves carbon emissions due to the energy required to capture, pump and transport the water. Detailed calculations for lawns are sparse. Townsend-Small & Czimczik (2010b) estimate up to $0.5 \text{ t ha}^{-1} \text{ y}^{-1}$ carbon emissions (10.6 tonnes of carbon per hectare over two decades). Studies of agricultural irrigation suggest similar numbers: $0.1\text{--}0.5 \text{ t C ha}^{-1} \text{ y}^{-1}$ (Griffiths-Sattenspiel & Wilson, 2009; Nelson *et al.*, 2009; Zou *et al.*, 2015).

Therefore, grass that is mown, fertilised and irrigated may produce emissions over two decades equal to as much as 43.9 tonnes of carbon equivalent per hectare. Some estimates of emissions due to lawn maintenance are as high as 126 tonnes carbon equivalent per hectare over two decades (Kong *et al.*, 2014) (Table 1). Such emission rates far outweigh the highest reported gross soil sequestration of 28 tonnes over the same timeframe. Beyond our two decade horizon, emissions from lawns will always outweigh initial soil carbon sequestration. Kong *et al.* (2014) estimated that the

carbon sink capacity of the lawns they studied would be offset by carbon emissions in 5–24 years under their current management, thereby shifting them from carbon sinks to permanent carbon sources. This number is within the range of data we gathered from multiple sources (Figure 1). Greenhouse gas emissions related to even basic lawn maintenance (i.e. infrequent mowing without fertiliser or irrigation) will eventually outweigh the carbon storage potential, transforming lawns from carbon sinks into carbon sources (Selhorst & Lal, 2013). Given that many lawns are old and have been in place for many decades, most of them can be assumed to act as carbon sources.

Finally, the carbon stored above ground in mown grass is 1.0 t ha^{-1} on average and 1.4 t ha^{-1} below ground (Guertal, 2012) (Table 2). This is negligible compared to the potential for carbon stored in plant tissue of trees and shrubs.

4. Carbon storage potential from converting lawns into treescapes

The reported carbon stored in plant tissue in natural forests varies considerably but can, for example, be as much as 690 t ha^{-1} in tropical Ghana (Nero *et al.*, 2017) and 360 t ha^{-1} in cool temperate climates such as New Zealand (Paul *et al.*, 2021) (Table 2). Even low shrub vegetation can store quantities of carbon that are significantly higher than those in lawns. For example, native sage scrub in California contains 43 t ha^{-1} of above and below ground carbon (Wheeler *et al.*, 2016). While these pristine natural forests and dense shrublands may not be compared directly to the carbon storage potential of urban treescapes, some report almost as high stores of carbon from parks: up to 420 t ha^{-1} of above ground carbon in tropical Ghana (475 t C ha^{-1} roots included) (Nero *et al.*, 2017) and up to 289 t ha^{-1} of above ground carbon in the cool temperate climate of Leicester, England (Davies *et al.*, 2011) (Table 2). Residential trees in Florida are reported to store 63 t of above ground carbon per hectare, and in Leipzig, Germany, above ground carbon stored per hectare of tree cover in areas with multi-story houses is as high as 64 tonnes (Strohbach & Haase, 2012).

Reported above and below ground sequestration rates of carbon average 61.2 t ha^{-1} over two decades in urban areas across the USA but over the same timeframe above ground carbon sequestration alone is as high as 137 t ha^{-1} in Seoul urban forest (Table 2) (Lee *et al.*, 2019; Nowak *et al.*, 2013). Shrubland can sequester up to 62 t ha^{-1} over two decades (Kimberley *et al.*, 2014). However, sequestration cannot continue indefinitely and approaches zero as rates of respiration match rates of photosynthesis. For example, net carbon flux does not differ significantly from zero in natural forests across New Zealand (Paul *et al.*, 2021). In an urban context, the fate of removed biomass will be important. If the wood from felled or pruned trees is used for furniture or to replace fossil fuels the carbon balance can remain positive.

In addition to the much higher carbon storage in above and below ground plant tissue in treescapes compared to lawns, the potential to store SOC must be considered. Soils under natural forests store more organic carbon than those under natural grasslands in the same climatic zones (Jobbágy & Jackson, 2000). Carbon content in soil under urban trees can be as high as 144 t ha^{-1} without the addition of fertiliser or irrigation (Dorendorf *et al.*, 2015) (Table 1). Furthermore, SOC under mown grass increases with the addition of trees or shrubs in a linear relationship with aboveground tree biomass (Bae & Ryu, 2015; Huyler

Table 1. A sample of publications reporting carbon equivalent emitted due to maintenance and organic carbon stored or sequestered in soil

Description	20-year emissions due to fertiliser use	20-year emissions due to fuel use for maintenance	20-year emissions due to irrigation	20-year gross organic carbon sequestration in soil	Net 20-year organic carbon sequestration in soil	Reported stored organic carbon in soil	Location	References
Mown lawns, low fertiliser rate (0–20 cm depth)	4.9	6.7	10.6	28.2	5.9		California	(Townsend-Small & Czimczik, 2010a, 2010b)
Mown lawns, high fertiliser rate (0–20 cm depth)	26.6	6.7	10.6	28.2	−15.6			
Mown lawns	1.3	3.8				45.8	USA Average	(Selhorst & Lal, 2013)
Residential lawns (0–50 cm depth)				3.0			Alabama	(Huyler et al., 2014a)
Weekly mown lawns		3.4					Massachusetts	(Lerman & Contosta, 2019)
Three-weekly mown lawns		1.4						
Residential lawn maintenance		2.8					Florida	(Horn et al., 2015)
Residential tree maintenance		0.04						
Grass (high fertiliser rate)	13.4							(Gu et al., 2015)
Grass, push mower		4.29					Singapore	(Velasco et al., 2021)
Parkland shrubs (0–60 cm depth)						91.5	Helsinki	(Lindén et al., 2020)
Parkland lawn (0–60 cm depth)						73.0		
Mixed forest soil (0–1 m depth)						101.9	Seoul	(Bae & Ryu, 2015)
Lawn (0–1 m depth)						37.4		
Urban forest (0–30 cm)						89	Auckland	(Weissert et al., 2016)
Lawn (0–30 cm) 3/4 sites irrigated and fertilised						108		
Lawn (0–40 cm) (100 years old) fertilised					5.96	29.8	Salt Lake City	(Smith et al., 2018)
Forest restoration 7–8 years old (0–100 cm)						37.9–82.9	New York City	(Downey et al., 2021)
Park lawn C (14yrs old) (0–15 cm)	126*					31.5	Shenzen and Hong Kong	(Kong et al., 2014)
Lawn (high sand content) (0–10 cm)				4.2		11.5	Texas	(Sapkota et al., 2020)
Dry urban forest (0–30 cm)						75.7	Hamburg	(Dorendorf et al., 2015)
Wet urban forest (0–30 cm)						144.3		

All units are tonnes carbon equivalent ha^{−1}. See Supplementary Information for an extended version of table.

*Emissions associated with mowing, fertilising, irrigation, and pesticide application.




Fig. 1. Cumulative carbon sequestration/emissions over time (log scale) for lawns and treescapes with the assumption of starting at zero carbon content both below and above ground at year 1. Note that carbon gains level off after about 50 years for grass, and after about 100 years for trees. Conversely, carbon losses associated with mowing, fertilising, irrigation and trimming are constant and quickly outweigh potential carbon gains in grass soils. The 'carbon compensation point' (where emissions equal sequestration) occurs as early as after a few years in lawns but may never occur in forest or treescapes. If older trees are removed and either used to replace fossil fuel, or as construction timber, then the carbon balance looks even more in favour of trees. Coloured bands represent a range of values sourced from Tables 1 and 2.

et al., 2014b, 2017; Lerman & Contosta, 2019). From sites in Seoul, Bae and Ryu (2015) report 37.4 t ha^{-1} of organic carbon in the top metre of lawn soil whereas carbon under urban forest was 2.4 times greater (89 t ha^{-1}) and that under mixed forest 2.7 times greater (101.9 t ha^{-1}). Lindén et al. (2020) found more soil carbon under shrubs than under lawns (91.5 and 73 t ha^{-1} , respectively, 0–60 cm depth) in Helsinki parkland. By contrast, in Auckland higher concentrations of carbon under mown grass than under urban forest have been reported (108 and 89 t ha^{-1} respectively) (Weissert et al., 2016). However, this difference was probably due to three out of the four mown grass sites being subject to fertilisation and irrigation, which in turn will have led to carbon emissions.

Emissions from treescapes due to trimming and other maintenance need to be considered as permanent components of their carbon balance. A study in Florida found carbon emissions due to maintaining trees of 0.04 t ha^{-1} projected over two decades, a rate of greenhouse gas emission related to maintenance approximately two orders of magnitude lower than for lawns (Horn et al., 2015) (Table 1, Figure 1). While urban trees provide carbon sequestration benefits this may be offset to some extent by their potential cost in water consumption, especially in arid or

semi-arid landscapes where water is already scarce (Dwyer et al., 1992). Direct comparisons between water requirements for mown grass and trees are few. However, in semi-arid Los Angeles, Litvak et al. (2014) found evapotranspiration of irrigated turfgrass was an order of magnitude higher than tree transpiration and in summer, evapotranspiration of the lawns with trees was lower than lawns without trees. Litvak et al. concluded that planting trees that partially shade irrigated urban lawns could be a water-saving measure in semi-arid environments. In Colorado, a significant negative relationship between water consumption and tree canopy was found, suggesting that homes with greater tree canopy area were associated with less water use (Rasmussen et al., 2021), and two study sites in Spain show consistently greater irrigation requirements for turf grass than for shrubs and trees (Hof & Wolf, 2014).

In summary, tree cover in any form or shape presents a much greater potential for sequestering and storing carbon, both above and below ground. The mid- to long-term effects of replacing lawns with treescapes are visualised in Figure 1, with the main advantage of trees being the much higher above ground storage, combined with the summative nature of emissions related to lawn maintenance. If urban trees that die, or need removal for

Table 2. A sample of publications reporting carbon stored and sequestered in urban vegetation: tonnes of carbon per hectare of vegetation cover and per hectare of land

Description	20-year carbon sequestration per hectare of cover	Stored carbon per hectare of cover	Stored carbon per hectare of land	Location	References
Natural forest (AG)			19.8	Barcelona	(Chaparro & Terradas, 2009)
Natural forest (AG and BG)			690.4	Kumasi	(Nero et al., 2017)
Natural forest (AG and BG plus litter)			144-360.5	Auckland	(Paul et al., 2021)
Public owned sites (AG)			288.6	Leicester	(Davies et al., 2011)
Urban trees (AG and BG)	61.2			Average across 50 states	(Nowak et al., 2013)
Urban Forest (AG)	136.8		63.19	Seoul	(Lee et al., 2019)
Natural urban forest AG and BG to 10 cm depth		263.04		New York	(Pregitzer et al., 2021)
Urban dry forest (AG and BG)			123.2	Hamburg	(Dorendorf et al., 2015)
Forest park (AG and BG)			262.4	Almada	(Mexia et al., 2018)
Park trees (AG and BG)			474.7	Kumasi	(Nero et al., 2017)
Domestic garden trees		28.6		Leipzig	(Strohbach & Haase, 2012)
Domestic garden trees (multi-story houses)		63.8			
Residential trees			63.0	Florida	(Timilsina, et al., 2014)
Mixed shrub species (AG)	62.2			New Zealand	(Kimberley et al., 2014)
Sage scrub (AG and BG to 10 cm depth)		43		California	(Wheeler et al., 2016)
Grass (AG)		1.0		Average reported from 2012 literature review	(Guertal, 2012)
Grass (BG)		1.4			

AG, above ground stems; BG, below ground roots.

See Supplementary Information for an extended version of table.

other reasons, are used for timber or to replace fossil fuels, a more favourable carbon balance than shown in [Figure 1](#) will result.

5. The upside of treescapes

Lawns have been promoted as providing ecosystem services including carbon sequestration (Velasco *et al.*, 2021). By contrast, we demonstrate that in bioregions capable of supporting trees, mown grass represents a degraded ecosystem in terms of carbon relative to the forested land that once occupied these sites. The literature we review clearly demonstrates that treescapes not only store considerably more carbon per unit surface area than mown grass, but they remain carbon sinks for much longer time periods. Even more important than the larger carbon pools and sequestration rates offered by treescapes are the inevitable and constant emissions due to lawns, mostly caused by mowing, fertilisation and irrigation. With the ubiquity of mown grassland (Milesi *et al.*, 2005), an opportunity exists to create alternative landscapes that not only store more carbon above ground, but also have a potential for greater below-ground storage. Although the role of urban trees in providing ecosystem services, particularly carbon sequestration, has been evidenced in many cities around the world, little work has been done to implement this knowledge into land use policies (Haase *et al.*, 2014).

Planting trees in urban environments, in addition to fixing carbon, can reduce pollution, cool summer air temperatures thereby reducing heat stress-related mortality (Manickathan *et al.*, 2018; Singh *et al.*, 2012; Zölch *et al.*, 2016), mitigate storm-water runoff (O'Sullivan *et al.*, 2017), and if species native to the region are used, enhance ecological restoration. Lawns contribute to homogeneity of the landscape, and generally lack biodiversity, particularly if mowing occurs often (Ignatieva & Hedblom, 2018).

There are also social and health benefits associated with converting mown grass to trees. Increasing tree cover is usually favourable to residents for aesthetic reasons (O'Sullivan *et al.*, 2017) as well as for higher perceived safety (Mouratidis, 2019). Significant positive associations between tree cover and self-reported health of residents have been found, whereas no such benefits to health were associated with grass (Reid *et al.*, 2017). Significant improvement in cardiovascular health parameters have been found among people walking among trees in contrast to no effect achieved from city-walking (Lee & Lee, 2014). One study conducted in Athens, Greece found that people had higher heart rates when running on routes without trees and that they felt more calm and experienced more joy running in a treescape or a seascape than running in treeless environments (Paraskevopoulou *et al.*, 2022).

Although the majority of people value the presence of trees (Camacho-Cervantes *et al.*, 2014; Lusk *et al.*, 2020), some people dislike them due to the shade they create. Shading is, and will continue to be, an important consideration for city planners. Perceptions and behaviour towards trees in urban areas is complex and multidimensional (Camacho-Cervantes *et al.*, 2014). However, despite a diversity of opinion, a literature review by Mullaney *et al.* (2015) found that residents consistently view street trees positively, and most believe that the benefits provided by trees significantly outweigh any detriments. It is not surprising that shade trees are valued more in warm climates than in cold climates. For example, 93% of respondents in Florence thought the city needed more trees to provide shaded cool places (Speak & Salbitano, 2021). Nonetheless, tree cover has been found to raise winter temperatures in cold climates (Edmondson *et al.*,

2016). Community perceptions of trees are important. However, negative attitudes towards trees by some members of the public should not outweigh the imperative to address global warming.

Lawns have a significant influence on landscapes, forming a component of most urban green spaces. They can foster a sense of well-being, and access to them is viewed by many people as very valuable to the extent that many people in the Western world view them as a compulsory element of urban landscapes (Ignatieva *et al.*, 2017). Lawns have become such a common feature of our living environments that it is difficult for some people to imagine alternatives (Hellner & Vilkenas, 2014). Generally people do not question their social, or aesthetic values, let alone their ecological consequences (Ignatieva *et al.*, 2017). We do not suggest that all lawns should be converted into treescapes but the imperative of addressing climate change is of such importance that we believe there should be an effort to convert as much as possible, and there may be greater acceptance of such change than land managers imagine. One study conducted in Warsaw found that a clear majority of park visitors thought conversion of lawns to tree plantations would increase the attractiveness of the parks' interiors (Sikorska *et al.*, 2020). Ultimately, the societal and psychological complexities of the introduction of trees needs to be addressed separately; here, we are focusing on the aspect of the net carbon balance.

To maximise sequestering atmospheric carbon, local and central governments need to: firstly, quantify the area of mown grass and identify the potential for carbon sequestration by conversion to treescapes. Improvements in data quality from satellite images and improved classification algorithms now allow for accurate mapping of mown grassland cover (Weng, 2012). Secondly, specific policies and plans to maximise this conversion should be introduced. We propose four types of urban and non-urban space that, prior to human occupation, would have supported forest but are now degraded to lawns: public parks and recreational reserves, highway verges, residential street berms and private gardens. The relevance and suitability of these types of spaces for conversion to treescapes will vary depending on the country or even region in question. To evaluate the practicality and socio-ecological acceptance of the type of conversion is beyond the scope of this study.

6. Public parks and recreational reserves

We propose widespread re-evaluation of park management plans with the view to identifying areas to be retired from mown grass. This will require a fundamental shift in what people perceive as desirable and usable in parks. However, the climate crisis is of such magnitude that all possible options must be explored, and action taken to reduce emissions and increase carbon sequestration wherever possible. Planting approaches will need to take account of the quantity of public traffic the areas receive. We therefore propose a range of planting options, a mixture of which might be introduced into any given park. These options range from a fully tiered forest structure (canopy and emergent trees with under storey shrubs and ground vegetation such as ferns, woody shrubs, grasses and/or herbs depending on the location), to open-plan treescapes with mostly closed canopies but with an open understorey and gaps to provide spaces for people to gather for picnics and other activities, to low-density treescapes with indigenous grass species that require little or no mowing ([Figure 2](#)). Shrubland can be used where an open vista is required.




Fig. 2. Left-hand side: potential impact on the global carbon cycle if one-third of urban lawns are converted into treescapes globally. The model calculation is based on an average of three different planting regimes: restoration of a full forest ecosystem, open-plan treescapes with light gaps and low-density treescapes. On the right-hand side, a model calculation for New Zealand highway verges is shown, assuming a highway network of 94,000 km (not including minor roads) a verge of 1–2 m and a mix of low- and high-density plantings. Over two decades, this conversion would result in carbon storage of 1.2–2.3 Mt C 20 y⁻¹.

7. Road verges and private gardens

Highway margins are routinely cleared of vegetation and maintained as mown grass due to a perception that it makes roads safer. On the one hand, trees in close proximity to road margins create crash hazards for drivers that lose control, but, on the other hand, they reduce driver stress, lower driving speeds and thereby reduce the occurrence and severity of crashes (Van Treese II et al., 2017). There is, therefore, a case to be made for converting roadside mown verges into shrubs and trees that can contribute to carbon sequestration and remove the need for mowing (Figure 2). In New Zealand, for example, with a population of approximately 5 million, there is 94,000 km of highway (NZTA). Assuming 1–2 m of mown grass on each side of the road, we estimate a total area of 18,800–37,600 ha of mown highway road margins. New Zealand native shrubs can store up to 62.2 t ha⁻¹ of above ground carbon after two decades of growth thereby potentially providing 1.2–2.3 Mt of above ground carbon storage over two decades (Figure 2) (Kimberley et al., 2014). Planting with trees could store more over longer timeframes. Similarly, front and back lawns and streetside berms of grass are common in many countries especially those influenced by colonisation (Ignatieva & Stewart, 2009). Some of these areas provide spaces for children to play or to access below-ground infrastructure, but many exist without a specific reason as a default setting that is seldom questioned.

Much of this grass could be converted into shrubs or trees (Figure 2).

8. Recommendations and conclusions

Central and local governments throughout the world should consider introducing policies to regulate or incentivise the conversion of treeless mown grass areas into shrublands and/or treescapes, eliminating mown grass wherever possible, and where circumstances dictate, trees with minimal grass should be retained. Such a process will reduce greenhouse gas emissions associated with maintaining mown grass and add stored carbon to the landscape. To estimate the global potential of carbon sequestration from converting mown grass into trees we use 61.2 t ha⁻¹ for whole tree carbon, which is the average across 50 states in the USA (Nowak et al., 2013) (Table 2). Global urban lawn area is estimated at 150,000–800,000 km² (Ignatieva & Hedblom, 2018). If one-third of the lawn in urban areas could be converted to tree cover, we estimate that 0.31–1.63 Gt of carbon could be sequestered over two decades. The estimate might be ambitious, but even one-tenth of these figures would be substantial. The rate of sequestration will accelerate in following decades, and SOC sequestration as well as emission savings due to the redundant lawn maintenance will be additional to this estimate.

Importantly, a large proportion of the lawns in scope are in the developed world, and their conversion to treescapes will contribute to a more equitable global land use (Creutzig et al., 2019). Finally, the many co-benefits of trees most importantly for human health and biodiversity make a strong case to reconsider urban land use.

Supplementary material. The supplementary material for this article can be found at <https://doi.org/10.1017/sus.2023.1>

Acknowledgements. Not applicable.

Author contributions. LNG wrote the first draft and edited the manuscript, SL designed the figures, contributed to writing and editing, BB contributed to writing and editing.

Financial support. Not applicable.

Conflict of interest. There are no conflicts of interest, and all data are in the Supplementary Materials File.

References

- Bae, J., & Ryu, Y. (2015). Land use and land cover changes explain spatial and temporal variations of the soil organic carbon stocks in a constructed urban park. *Landscape and Urban Planning*, 136, 57–67. doi: <https://doi.org/10.1016/j.landurbplan.2014.11.015>.
- Camacho-Cervantes, M., Schondube, J. E., Castillo, A., & MacGregor-Fors, I. (2014). How do people perceive urban trees? Assessing likes and dislikes in relation to the trees of a city. *Urban Ecosystems*, 17, 761–773.
- Chaparro, L., & Terradas, J. (2009). Ecological services of urban forest in Barcelona. *Institut Municipal de Parcs i Jardins Ajuntament de Barcelona, Àrea de Medi Ambient*.
- Creutzig, F., d'Amour, C. B., Weddige, U., Fuss, S., Beringer, T., Gläser, A., Kalkuhl, M., Steckel, J. C., Radebach, A., & Edenhofer, O. (2019). Assessing human and environmental pressures of global land-use change 2000–2010. *Global Sustainability*, 2, E1. doi: <https://doi.org/10.1017/sus.2018.15>.
- Davies, Z. G., Edmondson, J. L., Heinemeyer, A., Leake, J. R., & Gaston, K. J. (2011). Mapping an urban ecosystem service: Quantifying above-ground carbon storage at a city-wide scale. *Journal of Applied Ecology*, 48(5), 1125–1134. doi: <https://doi.org/10.1111/j.1365-2664.2011.02021.x>.
- Dorendorf, J., Eschenbach, A., Schmidt, K., & Jensen, K. (2015). Both tree and soil carbon need to be quantified for carbon assessments of cities. *Urban Forestry & Urban Greening*, 14(3), 447–455. doi: <https://doi.org/10.1016/j.ufug.2015.04.005>.
- Downey, A. E., Groffman, P. M., Mejía, G. A., Cook, E. M., Sritrairat, S., Karty, R., & McPhearson, T. (2021). Soil carbon sequestration in urban afforestation sites in New York City. *Urban Forestry & Urban Greening*, 65, 127342. doi: <https://doi.org/10.1016/j.ufug.2021.127342>.
- Dwyer, J. F., McPherson, E. G., Schroeder, H. W., & Rowntree, R. A. (1992). Assessing the benefits and costs of the urban forest. *Journal of Arboriculture*, 18(5), 227–234. doi: [18:227-234](https://doi.org/10.1016/j.ufug.2015.04.005).
- Edmondson, J. L., Stott, I., Davies, Z. G., Gaston, K. J., & Leake, J. R. (2016). Soil surface temperatures reveal moderation of the urban heat island effect by trees and shrubs. *Scientific Reports*, 6(1), 33708. doi: [10.1038/srep33708](https://doi.org/10.1038/srep33708).
- Gilbert, O. (1989). *Gardens*. Springer.
- Griffiths-Sattenspiel, B., & Wilson, W. (2009). *The carbon footprint of water*. River Network, Portland.
- Gu, C., Crane, J., Hornberger, G., & Carrico, A. (2015). The effects of household management practices on the global warming potential of urban lawns. *Journal of Environmental Management*, 151, 233–242. doi: <https://doi.org/10.1016/j.jenvman.2015.01.008>.
- Guertel, E. A. (2012). Carbon sequestration in turf landscapes: A review. In R. Lal, & B. Augustin (Eds.), *Carbon sequestration in urban ecosystems* (pp. 197–213). Dordrecht, Netherlands: Springer.
- Haase, D., Larondelle, N., Andersson, E., Artmann, M., Borgström, S., Breuste, J., Gomez-Baggethun, E., Gren, Å., Hamstead, Z., Hansen, R., Kabisch, N., Kremer, P., Langemeyer, J., Rall, E. L., McPhearson, T., Pauleit, S., Qureshi, S., Schwarz, N., Voigt, A., ... Elmqvist, T. (2014). A quantitative review of urban ecosystem service assessments: Concepts, models, and implementation. *AMBIO*, 43(4), 413–433. doi: [10.1007/s13280-014-0504-0](https://doi.org/10.1007/s13280-014-0504-0).
- Hellner, A., & Vilkėnas, J. (2014). In search for sustainable alternatives to lawns—connecting research with landscape design. Master's Thesis. Swedish University of Agricultural Sciences, Uppsala.
- Hof, A., & Wolf, N. (2014). Estimating potential outdoor water consumption in private urban landscapes by coupling high-resolution image analysis, irrigation water needs and evaporation estimation in Spain. *Landscape and Urban Planning*, 123, 61–72.
- Horn, J., Escobedo, F. J., Hinkle, R., Hostetler, M., & Timilsina, N. (2015). The role of composition, invasives, and maintenance emissions on urban forest carbon stocks. *Environmental Management*, 55(2), 431–442. doi: [10.1007/s00267-014-0400-1](https://doi.org/10.1007/s00267-014-0400-1).
- Huyler, A., Chappelka, A. H., Fan, Z., & Prior, S. A. (2017). A comparison of soil carbon dynamics in residential yards with and without trees. *Urban Ecosystems*, 20(1), 87–96. doi: [10.1007/s11252-016-0572-y](https://doi.org/10.1007/s11252-016-0572-y).
- Huyler, A., Chappelka, A. H., Prior, S. A., & Somers, G. L. (2014a). Drivers of soil carbon in residential 'pure lawns' in Auburn, Alabama. *Urban Ecosystems*, 17(1), 205–219. doi: [10.1007/s11252-013-0294-3](https://doi.org/10.1007/s11252-013-0294-3).
- Huyler, A., Chappelka, A. H., Prior, S. A., & Somers, G. L. (2014b). Influence of aboveground tree biomass, home age, and yard maintenance on soil carbon levels in residential yards. *Urban Ecosystems*, 17(3), 787–805. doi: [10.1007/s11252-014-0350-7](https://doi.org/10.1007/s11252-014-0350-7).
- Ignatieva, M., Eriksson, F., Eriksson, T., Berg, P., & Hedblom, M. (2017). The lawn as a social and cultural phenomenon in Sweden. *Urban Forestry & Urban Greening*, 21, 213–223.
- Ignatieva, M., & Hedblom, M. (2018). An alternative urban green carpet. *Science*, 362(6411), 148–149. doi: [10.1126/science.aau6974](https://doi.org/10.1126/science.aau6974).
- Ignatieva, M. E., & Stewart, G. H. (2009). Homogeneity of urban biotopes and similarity of landscape design language in former colonial cities. In M. J. McDonnell, A. K. Hahs, & J. H. Breuste (Eds.), *Ecology of cities and towns: A comparative approach* (pp. 399–421). Cambridge University Press.
- Jobbágy, E. G., & Jackson, R. B. (2000). The vertical distribution of soil organic carbon and its relation to climate and vegetation. *Ecological Applications*, 10(2), 423–436. doi: [https://doi.org/10.1890/1051-0761\(2000\)010\[0423:TVDSOJ\]2.0.CO;2](https://doi.org/10.1890/1051-0761(2000)010[0423:TVDSOJ]2.0.CO;2).
- Kimberley, M., Bergin, D., & Beets, P. (2014). Technical article No. 10; carbon sequestration by planted native tree shrubs. In: Tane's Tree Trust.
- Kong, L., Shi, Z., & Chu, L. M. (2014). Carbon emission and sequestration of urban turfgrass systems in Hong Kong. *Science of the Total Environment*, 473–474, 132–138. doi: <https://doi.org/10.1016/j.scitotenv.2013.12.012>.
- Lee, D.-H., Kil, S.-H., Jo, H.-K., & Choi, B. (2019). Spatial distributions of carbon storage and uptake of urban forests in Seoul, South Korea. *Sensors and Materials*, 31(11), 3811–3826.
- Lee, J.-Y., & Lee, D.-C. (2014). Cardiac and pulmonary benefits of forest walking versus city walking in elderly women: A randomised, controlled, open-label trial. *European Journal of Integrative Medicine*, 6(1), 5–11. doi: <https://doi.org/10.1016/j.eujim.2013.10.006>.
- Lerman, S. B., & Contosta, A. R. (2019). Lawn mowing frequency and its effects on biogenic and anthropogenic carbon dioxide emissions. *Landscape and Urban Planning*, 182, 114–123. doi: <https://doi.org/10.1016/j.landurbplan.2018.10.016>.
- Lindén, L., Riikonen, A., Setälä, H., & Yli-Pelkonen, V. (2020). Quantifying carbon stocks in urban parks under cold climate conditions. *Urban Forestry & Urban Greening*, 49, 126633. doi: <https://doi.org/10.1016/j.ufug.2020.126633>.
- Litvak, E., Bijoor, N. S., & Pataki, D. E. (2014). Adding trees to irrigated turfgrass lawns may be a water-saving measure in semi-arid environments. *Ecophysiology*, 7, 1314–1330.
- Lusk, A. C., da Silva Filho, D. F., & Dobbett, L. (2020). Pedestrian and cyclist preferences for tree locations by sidewalks and cycle tracks and associated benefits: Worldwide implications from a study in Boston, MA. *Cities*, 106, 102111.
- Manickathan, L., Defraeye, T., Allegrini, J., Derome, D., & Carmeliet, J. (2018). Parametric study of the influence of environmental factors and tree properties on the transpirative cooling effect of trees. *Agricultural and Forest Meteorology*, 248, 259–274.

- Mexia, T., Vieira, J., Príncipe, A., Anjos, A., Silva, P., Lopes, N., & Pinho, P. (2018). Ecosystem services: Urban parks under a magnifying glass. *Environmental Research*, 160, 469–478. doi: <https://doi.org/10.1016/j.envres.2017.10.023>.
- Milesi, C., Running, S. W., Elvidge, C. D., Dietz, J. B., Tuttle, B. T., & Nemani, R. R. (2005). Mapping and modeling the biogeochemical cycling of turf grasses in the United States. *Environmental Management*, 36(3), 426–438. doi: 10.1007/s00267-004-0316-2
- Mouratidis, K. (2019). The impact of urban tree cover on perceived safety. *Urban Forestry & Urban Greening*, 44, 126434. doi: <https://doi.org/10.1016/j.ufug.2019.126434>.
- Mullaney, J., Lucke, T., & Trueman, S. J. (2015). A review of benefits and challenges in growing street trees in paved urban environments. *Landscape and Urban Planning*, 134, 157–166.
- Nelson, G. C., Robertson, R., Msangi, S., Zhu, T., Liao, X., & Jawar, P. (2009). *Greenhouse gas mitigation: Issues for Indian agriculture*: Intl Food Policy Res Inst.
- Nero, B. F., Callo-Concha, D., Anning, A., & Denich, M. (2017). Urban green spaces enhance climate change mitigation in cities of the global south: The case of Kumasi, Ghana. *Procedia Engineering*, 198, 69–83. doi: <https://doi.org/10.1016/j.proeng.2017.07.074>.
- Nowak, D. J., Greenfield, E. J., Hoehn, R. E., & Lapoint, E. (2013). Carbon storage and sequestration by trees in urban and community areas of the United States. *Environmental Pollution*, 178, 229–236. doi: <https://doi.org/10.1016/j.envpol.2013.03.019>.
- NZTA. Research and data. Retrieved from <https://www.nzta.govt.nz/roads-and-rail/research-and-data>.
- O'Sullivan, O. S., Holt, A. R., Warren, P. H., & Evans, K. L. (2017). Optimising UK urban road verge contributions to biodiversity and ecosystem services with cost-effective management. *Journal of Environmental Management*, 191, 162–171. doi: <https://doi.org/10.1016/j.jenvman.2016.12.062>.
- Paraskevopoulou, A. T., Chletsou, M., & Malesios, C. (2022). Runners experience lower heart rate, increased speed, and joy/calm on routes with trees, by the sea and through parks: Implications for climate change design. *Sustainability*, 14, 16280.
- Paul, T., Kimberley, M. O., & Beets, P. N. (2021). Natural forests in New Zealand – a large terrestrial carbon pool in a national state of equilibrium. *Forest Ecosystems*, 8(1), 34. doi: 10.1186/s40663-021-00312-0
- Pregitzer, C. C., Hanna, C., Charlop-Powers, S., & Bradford, M. A. (2021). Estimating carbon storage in urban forests of New York City. *Urban Ecosystems*, 25, 1–15.
- Qian, Y. L., Bandaranayake, W., Parton, W. J., Mechem, B., Harivandi, M. A., & Mosier, A. R. (2003). Long-term effects of clipping and nitrogen management in turfgrass on soil organic carbon and nitrogen dynamics. *Journal of Environmental Quality*, 32(5), 1694–1700. doi: <https://doi.org/10.2134/jeq2003.1694>.
- Rasmussen, S., Warziniack, T., Neel, A., O'Neil-Dunne, J., & McHale, M. (2021). When small is not beautiful: The unexpected impacts of trees and parcel size on metered water-use in a semi-arid city. *Remote Sensing*, 13, 998.
- Reid, C. E., Clougherty, J. E., Shmool, J. L. C., & Kubzansky, L. D. (2017). Is all urban green space the same? A comparison of the health benefits of trees and grass in New York City. *International Journal of Environmental Research and Public Health*, 14(11), 1411. Retrieved from <https://www.mdpi.com/1660-4601/14/11/1411>.
- Sapkota, M., Young, J., Coldren, C., Slaughter, L., & Longing, S. (2020). Soil physiochemical properties and carbon sequestration of urban landscapes in Lubbock, TX, USA. *Urban Forestry & Urban Greening*, 56, 126847. doi: <https://doi.org/10.1016/j.ufug.2020.126847>.
- Selhorst, A., & Lal, R. (2013). Net carbon sequestration potential and emissions in home lawn turfgrasses of the United States. *Environmental Management*, 51(1), 198–208. doi: 10.1007/s00267-012-9967-6
- Sikorska, D., Macegoniuk, S., Łaszkiewicz, E., & Sikorski, P. (2020). Energy crops in urban parks as a promising alternative to traditional lawns – perceptions and a cost-benefit analysis. *Urban Forestry & Urban Greening*, 49, 126579.
- Singh, D., Takahashi, K., Kim, M., Chun, J., & Adams, J. M. (2012). A hump-backed trend in bacterial diversity with elevation on Mount Fuji, Japan. *Microbial Ecology*, 63(2), 429–437.
- Smith, R. M., Williamson, J. C., Pataki, D. E., Ehleringer, J., & Dennison, P. (2018). Soil carbon and nitrogen accumulation in residential lawns of the Salt Lake Valley, Utah. *Oecologia*, 187(4), 1107–1118. doi: 10.1007/s00442-018-4194-3
- Speak, A. F., & Salbitano, F. (2021). Thermal comfort and perceptions of the ecosystem services and disservices of urban trees in Florence. *Forests*, 12, 1387.
- Strobbach, M. W., & Haase, D. (2012). Above-ground carbon storage by urban trees in Leipzig, Germany: Analysis of patterns in a European city. *Landscape and Urban Planning*, 104(1), 95–104. doi: <https://doi.org/10.1016/j.landurbplan.2011.10.001>.
- Timilsina, N., Staudhammer, C. L., Escobedo, F. J., & Lawrence, A. (2014). Tree biomass, wood waste yield, and carbon storage changes in an urban forest. *Landscape and Urban Planning*, 127, 18–27.
- Townsend-Small, A., & Czimczik, C. I. (2010a). Carbon sequestration and greenhouse gas emissions in urban turf. *Geophysical Research Letters*, 37(2), 1–5. <https://doi.org/10.1029/2009GL041675>.
- Townsend-Small, A., & Czimczik, C. I. (2010b). Correction to 'carbon sequestration and greenhouse gas emissions in urban turf'. *Geophysical Research Letters*, 37, L06707. doi: 10.1029/2010gl042735
- Van Treese II, J. W., Koeser, A. K., Fitzpatrick, G. E., Olexa, M. T., & Allen, E. J. (2017). A review of the impact of roadway vegetation on drivers' health and well-being and the risks associated with single-vehicle crashes. *Arboricultural Journal*, 39(3), 179–193. doi: 10.1080/03071375.2017.1374591
- Vasenev, V., & Kuzyakov, Y. (2018). Urban soils as hot spots of anthropogenic carbon accumulation: Review of stocks, mechanisms and driving factors. *Land Degradation & Development*, 29(6), 1607–1622.
- Velasco, E., Segovia, E., Choong, A. M. F., Lim, B. K. Y., & Vargas, R. (2021). Carbon dioxide dynamics in a residential lawn of a tropical city. *Journal of Environmental Management*, 280, 111752. doi: <https://doi.org/10.1016/j.jenvman.2020.111752>.
- Weissert, L. F., Salmond, J. A., & Schwendenmann, L. (2016). Variability of soil organic carbon stocks and soil CO₂ efflux across urban land use and soil cover types. *Geoderma*, 271, 80–90. doi: <https://doi.org/10.1016/j.geoderma.2016.02.014>.
- Weng, Q. (2012). Remote sensing of impervious surfaces in the urban areas: Requirements, methods, and trends. *Remote Sensing of Environment*, 117, 34–49.
- Wheeler, M. M., Dipman, M. M., Adams, T. A., Ruina, A. V., Robins, C. R., & Meyer, W. M. (2016). Carbon and nitrogen storage in California sage scrub and non-native grassland habitats. *Journal of Arid Environments*, 129, 119–125. doi: <https://doi.org/10.1016/j.jaridenv.2016.02.013>.
- Zölch, T., Maderspacher, J., Wamsler, C., & Pauleit, S. (2016). Using green infrastructure for urban climate-proofing: An evaluation of heat mitigation measures at the micro-scale. *Urban Forestry & Urban Greening*, 20, 305–316. doi: <https://doi.org/10.1016/j.ufug.2016.09.011>.
- Zou, X., Li, Y. E., Li, K., Cremades, R., Gao, Q., Wan, Y., & Qin, X. (2015). Greenhouse gas emissions from agricultural irrigation in China. *Mitigation and Adaptation Strategies for Global Change*, 20(2), 295–315.